arXiv:1108.2801v2 [math.DS] 27 Sep 2011

THE FIXED POINT OF THE PARABOLIC RENORMALIZATION OPERATOR

OSCAR LANFORD III, MICHAEL YAMPOLSKY

1. Introduction

In a recent paper [IS], H. Inou and M. Shishikura demonstrated that the successive parabolic renormalizations $\mathcal{P}^n(f_0)$ of the quadratic polynomial $f_0(z) = z + z^2$ converge to an analytic map f_* defined in a neighborhood of the origin, which satisfies the fixed point equation

$$\mathcal{P}(f_*) = f_*.$$

Conjecturally, the solution of the above functional equation is unique in a suitably restricted class of maps.

In this paper we present a class of analytic maps $\mathbf{P}$ which have a maximal analytic extension to a Jordan domain, satisfying the invariance property

$$\mathcal{P} : \mathbf{P} \to \mathbf{P}.$$

The covering properties of a map $f \in \mathbf{P}$ admit an explicit topological model. We prove that the Inou-Shishikura fixed point f_* of $\mathcal{P}$ is contained in $\mathbf{P}$, and conjecture that successive renormalizations of any map $f \in \mathbf{P}$ converge to f_*.

The boundary of the maximal domain of analyticity of f_* has a a highly degenerate geometry. It is this bad geometry that makes the study of f_* so challenging. In contrast, consider the parabolic renormalization of critical circle maps with a parabolic fixed point on the circle, which both of the authors have studied [Lan, Ya, EY]. The corresponding renormalization fixed point also has a maximal analytic extension, whose covering properties are similar to those of f_*. The geometry of its domain of analyticity, however, is rather tame, which permits both a numerical and an analytic study.

We present a numerical method for computing the Taylor's expansion of f_* with a high accuracy. Our approach also allows us to compute the domain $\mathrm{Dom}(f_*)$, and the reader will see the first computer-generated images of it. Finally, we obtain a numerical estimate of the leading eigenvalue of $D\mathcal{P}|_{f_*}$.

2. Local dynamics of a parabolic germ

2.1. **Fatou coordinates.** We briefly review the local dynamics an analytic function f in the vicinity of a parabolic fixed point at 0:

$$f(z) = e^{2\pi ip/q}z + O(z^2).$$

We consider first the case $q = 1$, that is, $f'(0) = 1$, and we write

$$f(z) = z + az^{n+1} + O(z^{n+2}), \tag{2.1}$$

for some $n \in \mathbb{N}$ and $a \neq 0$. The integer $n + 1$ can be recognized as the *multiplicity* of 0 as the solution of $f(z) - z = 0$.

A complex number ν of modulus one is called an *attracting direction* if the product $a\nu^n < 0$, and a *repelling direction* if the same product is positive. The terminology has the following meaning:

Proposition 2.1. *Let $\{f^k(z)\}$ be an orbit in $\mathrm{Dom}(f) \setminus \{0\}$ which converges to the parabolic fixed point 0. Then the sequence of unit vectors $f^k(z)/|f^k(z)|$ converges as $k \to \infty$ to one of the attracting directions.*

We say in this case that the orbit converges to p *from the direction of ν*.

If f has a parabolic fixed point at 0, it admits a *local inverse* there, by which we mean a function g, defined and analytic on a neighborhood of 0, so that $g(f(z)) = z = f(g(z))$ for z near enough to 0. The germ at 0 of a local inverse is unique, but its domain of definition typically has to be chosen. A local inverse also has a parabolic fixed point at 0; attracting directions for f are repelling for the inverse and vice versa.

Definition 2.1. Let ν be an attracting direction for f. An *attracting petal for f (from the direction ν)* is a Jordan domain P with closure in $\mathrm{Dom}(f)$ such that:

1. $0 \in \partial P$
2. f is injective on P;
3. $f(\overline{P} \setminus \{0\}) \subset P$;
4. for any $z \in P$, the orbit $f^k(z)$ converges to 0 from the direction ν, and the convergence of f^k to 0 is uniform on P.
5. conversely, any orbit $f^k(z)$ which converges to z from the direction ν is eventually in P.

Similarly, U is a *repelling petal* for f if it is an attracting petal for some local inverse g of f.

Judiciously chosen petals can be organized into a *Leau-Fatou Flower* at 0:

Theorem 2.2. *There exists a collection of* n *attracting petals* P_i^a*, and* n *repelling petals* P_j^r *such that the following holds. Any two repelling petals do not intersect, and every repelling petal intersects exactly two attracting petals. Similar properties hold for attracting petals. The union*

$$(\cup P_i^a) \cup (\cup P_j^r) \cup \{0\}$$

forms an open simply-connected neighborhood of 0*.*

The proof of this statement relies on some changes of coordinates. First of all: Every germ of the form (2.1) can be brought into the form

$$f(z) = z + z^{n+1} + \alpha z^{2n+1} + O(z^{3n+1}) \tag{2.2}$$

in a suitable conformal local coordinate change at 0. In fact, a straightforward induction shows the following:

Proposition 2.3 (cf. [Mil1] Problem 10-d,[BE])**.** *For every germ of the form (2.1) there exists a unique* $\alpha \in \mathbb{C}$ *such that for every* $N \in \mathbb{N}$ *greater that* $2n + 1$ *there is a locally conformal change of coordinates* ψ*, with* $\psi(0) = 0$*, such that*

$$\psi \circ f \circ \psi^{-1}(z) = z + z^{n+1} + \alpha z^{2n+1} + O(z^N).$$

Further, there exists a formal power series $\Psi(z) = \sum_{k=1}^{\infty} p_k z^k$ *which formally conjugates*

$$\Psi \circ f \circ \Psi^{-1}(z) = z + z^{n+1} + \alpha z^{2n+1}.$$

Thus, the number $\alpha \in \mathbb{C}$ *is a formal conjugacy invariant of* f*, and specifies its formal conjugacy class uniquely.*

For the next few paragraphs, we will take f to have the special form (2.2). The attracting directions are then the nth roots of -1. We will describe some ways of constructiong attracting petals for the attracting direction ν; the adjustments necessary to deal with repelling petals are routine. The reader is reminded that (2.2) is *not* the general form of a mapping with a parabolic fixed point of order n; it has been cleaned up by making a preliminary analytic change of coordinates to eliminate some powers of z in its Taylor series.

The behavior of orbits of such an f near 0 is greatly clarified by making the coordinate change

$$w = \kappa(z) := -\frac{1}{nz^n}.$$

We are considering a particular attracting direction, and we take κ to be defined on the sector between the two adjacent repelling directions; it opens up this sector to the complex plane cut along the positive real axis. With its domain of definition restricted in this way, κ is bijective, and its inverse is given by

$$\kappa^{-1}(w) = \left(-\frac{1}{nw}\right)^{1/n}$$

where the branch of the n-th root is the one cut along the positive real axis and taking the value ν at -1.

Then

$$\begin{aligned}
F(w) &:= \kappa \circ f \circ \kappa^{-1}(w) \\
&= -\frac{1}{n}\left(f\left(\left(-\frac{1}{nw}\right)^{\frac{1}{n}}\right)\right)^{-n} = \\
&= -\frac{1}{n}\left(\left(-\frac{1}{nw}\right)^{\frac{1}{n}} + \left(-\frac{1}{nw}\right)^{\frac{n+1}{n}} + \alpha\left(-\frac{1}{nw}\right)^{\frac{2n+1}{n}} + O\left(\left(\frac{-1}{nw}\right)^{\frac{3n+1}{n}}\right)\right)^{-n} = \\
&= w \cdot \left(1 + \left(-\frac{1}{nw}\right) + \alpha\left(\frac{1}{nw}\right)^2 + O\left(\frac{1}{w^3}\right)\right)^{-n} = \\
&= w \cdot \left(1 + \frac{1}{w} - \frac{\alpha}{n} \cdot \frac{1}{w^2} + \frac{n+1}{2n} \cdot \frac{1}{w^2} + O(\frac{1}{w^3})\right).
\end{aligned}$$

We thus obtain:

$$F(w) = w + 1 + \frac{A}{w} + O(1/w^2) \quad \text{as } w \to \infty$$

where

$$A = \frac{1}{n}\left(\frac{n+1}{2} - \alpha\right).$$

Selecting a right half-plane $H_r = \{\operatorname{Re} z > r\}$ for a sufficiently large $r > 0$, we have

$$\operatorname{Re} F(w) > \operatorname{Re} w + 1/2, \text{ and hence } F(\overline{H_r}) \subset H_r.$$

The domain $\kappa^{-1}(H_r)$ is then an attracting petal for the attracting direction ν. In the case of a simple parabolic point, what we have just shown simplifies to the assertion that any disk of sufficiently small radius tangent to the imaginary axis from the left at 0 is an attracting petal.

The petals just discussed – pullbacks of half-planes under κ – have boundaries tangent at the origin to the directions $e^{\pm i\pi/(2n)}\nu$. For many purposes – such as the proof of Theorem 2.2 – we will need petals with strictly larger opening angle. There are many ways to construct such petals; here is one which is convenient for our purposes. Let $\pi/2 < \alpha < \pi$, $R > 0$, and let

$$\Delta(\alpha, R) := \{w : -\alpha < \operatorname{Arg}(w - R) < \alpha\}$$

(i.e., $\Delta(\alpha, R)$ is the sector $\{-\alpha < \operatorname{Arg}(w) < \alpha\}$ translated right by R.) From

$$F(w) = w + 1 + O(1/w),$$

there exists a $R_0 = R_0(\alpha)$ so that

$$|\operatorname{Arg}(F(w) - w)| < \pi - \alpha \quad \text{and} \quad \operatorname{Re}(F(w)) > \operatorname{Re}(w) + 1/2 \tag{2.3}$$

for $|w| \geq R_0$. If R is large enough, the domain $\Delta(\alpha, R)$ does not intersect the disk $\{|w| \leq R_0\}$, so (2.3) holds for $w \in \Delta(\alpha, R)$. For such R's, by elementary geometric considerations,

$$F\left(\overline{\Delta(\alpha, R)}\right) \subset \Delta(\alpha, R),$$

and $F^n(w) \underset{n\to\infty}{\longrightarrow} \infty$ for all $w \in \Delta(\alpha, R)$. Further any $\Delta(\alpha, R)$ contains a right half-plane and hence eventually contains any F-orbit converging to ∞. Finally, it can be verified that the sequence of iterates F^n converges uniformly to ∞ on $\Delta(\alpha, R)$. We omit this verification; it uses simplified versions of the ideas used in the proof of Lemma 2.13. Thus, sets of the form $\kappa^{-1}(\Delta(\alpha, R))$ are attracting petals, symmetric about the attracting direction under consideration, with tengents at the origin in directions $e^{\pm i\alpha/n}\nu$. It will be useful to have a general term for behavior for this: We will say that a petal P with attracting or repelling direction ν is *ample* if

$$P \supset \{z : |\operatorname{Arg}(z/\nu)| < \alpha/n, |z| < r\}$$

for some $\alpha > \pi/2$ and sufficiently small r.

The dynamics inside a petal is described by the following:

Proposition 2.4. *Let P be an attracting petal for f. Then there exists a conformal change of coordinates ϕ defined on P, conjugating $f(z)$ to the unit translation $T : z \mapsto z + 1$.*

Proof. For a traditional proof, see e.g. [Mil1] §10. We cannot resist giving a proof based on quasiconformal surgery, which probably originated in the work of Voronin [Vor]. For definiteness, we discuss the case of an attracting petal with attracting direction ν, and let

$$F(w) = w + 1 + O(1/w)$$

be as above. Also as above, we select a right half-plane H_r. The main step will be to prove the existence of ϕ for the special petal $\kappa^{-1}H_r$, which we provisionally denote by P_0. The case of a general petal will then follow by an easy extension argument.

As we know, $F(\overline{H_r}) \subset H_r$, so let us denote S the closed strip

$$S = \overline{H_r \setminus F(H_r)}.$$

Setting $\mathbb{S} = \{\operatorname{Re} z \in [0,1]\}$, let h be any diffeomorphism

$$h : S \to \mathbb{S},$$

which on the boundary of the strip conjugates F to T:

$$T \circ h(w) = h \circ F(w), \text{ for all } w \text{ with } \operatorname{Re} w = r.$$

We will further require that the first partial derivatives of h and h^{-1} be uniformly bounded in S. Verifying the existence a diffeomorphism with these properties is an elementary exercise which we leave to the reader.

The diffeomorphism h defines a new complex structure $\mu = h_*\sigma_0$ on $\mathbb{S}$, which we extend to the left half-plane $\{\operatorname{Re} z < 0\}$ by

$$\mu|_w = (T^n)^*\mu \text{ for } T^n(w) \in \mathbb{S}.$$

Gluing together H_r with the standard complex structure and the half-plane $\{\operatorname{Re} z < 1\}$ with structure μ via the homeomorphism h (which is now analytic), and using the Measurable Riemann Mapping Theorem, we obtain a new Riemann surface X. By the Uniformization Theorem, X is conformally isomorphic either to $\mathbb{C}$ or to the disk. By construction, X is quasiconformally isomorphic to $\mathbb{C}$ and therefore cannot be conformally isomorphic to the disk. We can specify a conformal isomorphism $\Phi : X \to \mathbb{C}$ uniquely by imposing normalization conditions $\Phi(0) = 0$ and $\Phi(-1) = -1$.

The pair of maps $T|_{\{\operatorname{Re} z<0\}}$ and $F|_{H_r}$ induces a conformal automorphism of X, which we denote by $\tilde{F}$. Then $\Phi \circ \tilde{F} \circ \Phi^{-1}$ is a conformal automorphism of $\mathbb{C}$ with no fixed point. It is a standard fact that the only such automorphisms are translations, and our choice of normalization for Φ implies that

$$\Phi \circ \tilde{F} \circ \Phi^{-1} \equiv T. \tag{2.4}$$

But $\tilde{F} = F$ on $H_r \subset X$, so we get

$$\Phi \circ F = T \circ \Phi \quad \text{on } H_r.$$

Moreover, the restriction of Φ to H_r is analytic in the standard sense. Thus, we set $\phi := \Phi \circ \kappa$ on $P_0 := \kappa^{-1} H_r$ and obtain

$$\phi \circ f = \phi + 1 \quad \text{on } P_0,$$

as desired. Since Φ is a conformal isomorphism from X to $\mathbb{C}$, the map ϕ is univalent on P_0.

This proves the existence of ϕ on the particular petal P_0. We provisionally denote the above ϕ, which is defined on P_0, by ϕ_0. We define

$$B_\nu^f := \{z : f^n(z) \to 0 \quad \text{from the direction } \nu\}.$$

If $z_0 \in B_\nu^f$, then $f^n(z_0) \in P_0$ for sufficiently large n. If $f^{n_0}(z_0) \in P_0$, then $(f^{n_0})^{-1}P_0$ is an open set containing z_0 and contained in B_ν^f, so B_ν^f is open. Since P_0 is mapped into itself by f, and since

$$\phi_0(f(z)) = \phi_0(z) + 1 \quad \text{on } P_0,$$

$\phi_0(f^n(z_0)) - n$ takes the same value for all n for which $f^n(z_0) \in P_0$. We denote this common value by $\phi(z_0)$, thus obtaining a function ϕ defined on all of B_ν^f and extending ϕ_0 defined on $P_0 \subset B_\nu^f$. Tautologically,

$$\phi(f(z)) = \phi(z) + 1.$$

If $f^{n_0}(z_0) \in P_0$, then $f^{n_0}(z) \in P_0$ on a neighborhood of z_0, so $\phi(z) = \phi_0(f^{n_0}(z)) - n_0$ on this neighborhood, which shows that the extended ϕ is analytic, but not necessarily univalent, on all of B_ν^f.

Now let P be a general attracting petal with the same attracting direction ν. By definition of petal, $P \subset B_\nu^f$, so we can restrict ϕ to P, thus obtaining an analytic function satisfying

$$\phi(f(z)) = \phi(z) + 1 \quad \text{on } P.$$

It remains to show that the restriction of ϕ to P is univalent. To see this, let z_1, z_2 be points of P with $\phi(z_1) = \phi(z_2)$. For sufficiently large n, $f^n(z_1)$ and $f^n(z_2)$ are both in P_0, so

$$\phi_0(f^n(z_1) = \phi(z_1) + n = \phi(z_2) + n = \phi_0(f^n(z_2)).$$

But, by construction, ϕ_0 is univalent, so $f^n(z_1) = f^n(z_2)$. The argument so far works for any pair z_1, z_2 in B_ν^f with $\phi(z_1) = \phi(z_2)$. Now, however, we use that facts that z_1 and z_2 are both in the petal P, which is mapped into itself by f and on which f is univalent. Hence, from $f^n(z_1) = f^n(z_2)$ it follows that $z_1 = z_2$, proving univalence of ϕ on P. □

We define *attracting Fatou coordinate* (for the attracting petal P with attracting direction ν) to be a function ϕ_A defined, analytic and univalent on P and satisfying

$$\phi_A(f(z)) = \phi_A(z) + 1 \quad \text{on } P.$$

As we have seen, such a function extends uniquely, via the above functional equation, to all of B_ν^f, and the extension restricts to an attracting Fatou coordinate on any other petal with attracting direction ν. It is clear that, if ϕ_A is an attracting Fatou coordinate then so is $\phi_A + c$ for any constant c. We will see shortly that any two attracting Fatou coordinates differ only in this way.

Any attracting Fatou coordinate can be written in the form $\phi_A = \Phi_A \circ \kappa$, where Φ_A satisfies the functional equation

$$\Phi_A(F(w)) = \Phi_A(w) + 1 \quad (\text{with } F = \kappa \circ f \circ \kappa^{-1} \text{ as above,})$$

on an appropriate F-invariant domain "near infinity". We will refer to such Φ_A's as *Fatou coordinates at infinity.*

Repelling Fatou coordinates are defined similarly. Roughly speaking, a repelling Fatou coordinate for f means an attracting Fatou coordinate for an analytic local inverse g of f. If, however, $f(z) = z + z^{n+1} + \ldots$, then $g(z) = z - z^{n+1} + \ldots$, which can be brought back into the standard form by conjugating with $z \mapsto -z$ (n odd) or $z \mapsto i \cdot z$ (n even.) The factors of -1 respectively i propagate to numerous places in the formulas. We leave it to the reader to sort this all out except to specify that

- a repelling Fatou coordinate ϕ_R satisfies *the same* functional equation
$$\phi_R(f(z)) = \phi_R(z) + 1$$
as does an attracting one, but the domains are different, and
- the image of a repelling Fatou coordinate is mapped into itself by the unit *left* translation $w \mapsto w - 1$ (so, as it turns out, the image of an ample repelling petal under a repelling Fatou coordinate contains a *left* half-plane.

Again, it is useful to consider also *repelling Fatou coordinates at infinity*: If ϕ_R is a repelling Fatou coordinate, the corresponding one at infinity is

$$\Phi_R(w) = \phi_R(\kappa^{-1}(w))$$

(but the appropriate branches of κ^{-1} are different from the ones in the attracting case.)

Our next step is to prove a crude asymptotic formula for a Fatou coordinate at infinity. It is advantageous here to deviate from what we have been doing. We consider a mapping f of the form

$$f(z) = z + z^{n+1} + f_{n+2}z^{n+2} + \cdots ,$$

i.e., we do not assume we have made a preliminary change of variable to eliminate, e.g., the terms z^j for j between $n + 1$ and $2n + 1$. We introduce $F = \kappa \circ f \circ \kappa^{-1}$ as before; this time, the behavior of F near infinity is

$$F(w) = w + v(w) \quad \text{where } v(w) = 1 + v_1 w^{-1/n} + v_2 w^{-2/n} + \ldots;$$

the series converges for sufficiently large $|w|$. Let Φ_A be an attracting Fatou coordinate at infinity. By what we have already proved: For any $\alpha < \pi$, there is an R so that Φ_A extends analytically to a univalent function on the set $\{|w| > R, -\alpha < \operatorname{Arg}(w) < \alpha\}$.

Proposition 2.5.

$$\Phi_A'(w) \to 1 \quad \textit{and} \quad \Phi_A(w)/w \to 1$$

uniformly as $w \to \infty$ *in any sector* $\{-\alpha < \operatorname{Arg}(w) < \alpha\}$ *with* $\alpha < \pi$. *The same limits hold for* Φ_R, *but with* $w \to \infty$ *in the opposite sector* $\{-\alpha < \operatorname{Arg}(-w) < \alpha\}$

This proposition is a less-precise version of Lemma A.2.4 of [Sh], and the argument we give is the first part of Shishikura's proof of that lemma. Shishikura carries the analysis further and is able to identify, in favorable cases, the first correction to the indicated asymptotic behaviors. We do not give his full argument here, as we will prove Theorem 2.12, which gives more precise information about the asymptotic behavior of Fatou coordinates.

Proof. Fix α with $\pi/2 < \alpha < \pi$, and let $\alpha < \alpha_1 < \pi$. Take R_1 so that Φ_A is defined and univalent in

$$S_1 := \{|w| > R_1, -\alpha_1 < \operatorname{Arg}(w) < \alpha_1\}$$

and also so that $|v(w) - 1| < 1/4$ on S_1. For $w_0 \in S_1$, denote by $\rho = \rho(w_0)$ the distance from w_0 to the boundary of S_1. We will investigate limits as $|w_0| \to \infty$ in the strictly smaller sector $-\alpha < \operatorname{Arg}(w_0) < \alpha$; then there is a constant $k > 0$ so that, asymptotically, $\rho(w_0) \geq k \cdot |w_0|$. In the following, we will frequently assume silently that $\rho(w_0)$ is "large enough". We will also use C to denote a generic "universal" constant; different instances of C need not denote the same constant.

For the first step, we use the Koebe Distortion Theorem: If $|w - w_0| < \rho - 2$ – so the disk of radius 2 about w is in $S_!$ – the mapping

$$a \mapsto \frac{\Phi_A(w+a) - \Phi_A(w)}{\Phi_A'(w)}$$

is analytic and univalent on $\{|a| < 2\}$ and has unit derivative at the origin. A simple rescaling of the Koebe Theorem to adapt it to the disk of radius 2 gives a universal constant $C > 0$ so that

$$\left| \frac{\Phi_A(w+a) - \Phi_A(w)}{\Phi_A'(w)} \right| > C^{-1} \quad \text{for } 3/4 < |a| < 5/4$$

We insert $a = v(w)$ into this estimate, use $|1 - v(w)| < 1/4$ to ensure that $3/4 < |v(w)| < 5/4$, and use also the functional equation

$$\Phi_A(w + v(w)) = \Phi_A(F(w)) = \Phi_A(w) + 1$$

to get

$$|\Phi'_A(w)| < C \quad \text{for } |w - w_0| < (\rho - 2).$$

Applying the Cauchy estimates gives a bound

$$|\Phi''_A(w)| \leq \frac{C}{\rho} \quad \text{for } |w - w_0| < \rho/2$$

(with a different C).

Next we apply Taylor's Formula with Integral Remainder to write

$$\Phi_A(w_0 + a) = \Phi_A(w_0) + a \cdot \Phi'(w_0) + a^2 \cdot \int_{s=0}^{1} (1 - s)\Phi''(w_0 + s \cdot a)ds.$$

Again, we set $a = v(w_0)$ and use $\Phi_A(w_0 + v(w_0)) = \Phi_A(w_0) + 1$ to get

$$1 - v(w_0)\Phi'_A(w_0) = (v(w_0))^2 \int_{s=0}^{1} (1 - s)\Phi''(w_0 + s \cdot a)ds.$$

Since the estimate $|\Phi''_A(w)| \leq C/\rho$ holds for all w appearing in the integral on the right, we get

$$|1 - v(w_0) \cdot \Phi'_A(w_0)| \leq C/\rho(w_0).$$

We have already remarked that $\rho(w_0) \geq k|w_0|$ as $w_0 \to \infty$ in the sector $\{-\alpha < \operatorname{Arg}(w_0) < \alpha\}$ so

$$|\Phi'_A(w_0) - v(w_0)^{-1}| = O(|w_0|^{-1}) \quad \text{in that sector.}$$

This establishes the asserted convergence of Φ'_A; the assertion about $\Phi(w)/w$ follows by integration.

$\square$

Equipped with this information about the asymptotic behavior of Fatou coordinates, we can now show that the image of a Fatou coordinate is large enough. As usual, it suffices – up to insertion of some minus signs – to consider the attracting case. Let Φ_A be an attracting Fatou coordinate, and let $0 < \alpha < \pi$. Then, for sufficiently large R, Φ_A extends analytically to a univalent function on

$$S := \{w : |w| > R, -\alpha < \operatorname{Arg}(w) < \alpha\}$$

Proposition 2.6. *Let $0 < \alpha_0 < \alpha$. Then, for sufficiently large R_0,*

$$\Phi_A(S) \supset S_0 := \{|w| < R_0, -\alpha_0 < \operatorname{Arg}(w) < \alpha_0\}.$$

Proof. Let $w_0 \in S_0$; we want to investigate solutions to the equation $\Phi_A(w) = w_0$ which we rewrite as

$$w = w_0 + w - \Phi_A(w) =: \Psi_0(w)$$

The idea is to apply the Contraction Mapping Principle to Ψ_0, using the fact that $\Psi_0'(w) = 1 - \Phi_A'(w)$ which is small for w large. To do this, we need to find a domain mapped into itself by Ψ_0 and on which Ψ_0 is contractive. Suppose we can find a $\delta > 0$ so that

- $|\Psi_0'(w)| < 1/2$ for $|w - w_0| < \delta$
- $|\Psi_0(w_0) - w_0| < \delta/2$

Then, for $|w - w_0| < \delta$,

$$\begin{aligned} |\Psi_0(w) - w_0| &\le |\Psi_0(w) - \Psi_0(w_0)| + |\Psi_0(w_0) - w_0| \\ &< (1/2) \cdot |w - w_0| + \delta/2 \le \delta/2 + \delta/2 = \delta, \end{aligned}$$

so the disk of radius δ about w_0 will be mapped to itself, and Ψ_0 will have a unique fixed point in this disk.

We implement this strategy as follows: First of all we arrange, by making R larger if necessary, that $|\Phi_A'(w) - 1| < 1/2$ on S. We write

$$\epsilon := \sin(\alpha - \alpha_0),$$

and we note that, by elementary geometry, $|w - w_0| < \epsilon \cdot w_0$ implies $|\operatorname{Arg}(w/w_0)| < \alpha - \alpha_0$. If we further take $R_0 \ge (1 - \epsilon)^{-1} R$, then the disk of radius $\delta := \epsilon \cdot |w_0|$ about w_0 is contained in S, for any $w_0 \in S_0$. Recall that we have already arranged that $|\Psi_0'| < 1/2$ on S. Finally, we apply Proposition 2.5 to see that, by taking R_0 large enough we can arrange that

$$|\Phi_A(w_0) - w_0| < (1/2) \cdot \epsilon \cdot |w_0| \quad \text{for all } w_0 \in S_0.$$

All the element for the above contraction argument are now in place, and we can conclude that, for every $w_0 \in S_0$, there is a unique w with $\Phi_A(w) = w_0$ in $\{|w - w_0| < \epsilon|w_0|\} \subset S$. This proves the assertion $\Phi_A(S) \supset S_0$.

□

It follows from this proposition that:

Proposition 2.7. *The image under ϕ_A of any ample petal of f contains a right half-plane.*

A similar assumption holds for ample repelling petals, but with the image under ϕ_R covering a left half-plane.

Let P be an attracting petal. We define a relation on P by $z_1 \sim z_2$ if z_1 and z_2 are on the same orbit, that is, if either $z_2 = f^j(z_1)$ or

$z_1 = f^j(z_2)$ (with $j \geq 0$.) It is easy to check that this is an equivalence relation.

Consider the quotient $P/_\sim$. The canonical projection $\pi : P \to P/_\sim$ is locally injective, and it is straightforward to verify that there is a unique way to give $P/_\sim$ a Riemann surface structure in such a way as to make π analytic and therefore a local conformal isomorphism.

Let P_1 be another attracting petal contained in P. Since the orbit of every point $z \in P$ eventually lands in P_1, the inclusion $P_1 \hookrightarrow P$ induces a conformal homeomorphism

$$P_1/\sim \underset{\simeq}{\longrightarrow} P/_\sim.$$

Now if P_2 is any attracting petal with the same attracting direction as P, the intersection $P \cap P_2$ is also a petal. Hence

$$P_2/_\sim \simeq P/_\sim \simeq (P \cap P_2)/_\sim.$$

Thus, the quotient $P_\sim$ does not depend on the choice of the petal P but only on the choice of the attracting direction ν corresponding to P. We will write

$$P/_\sim \equiv \mathcal{C}^\nu_A.$$

We will omit ν from the notation when the choice of the attracting direction is clear from the context (for instance, when there is only one attracting direction).

Let ϕ_A be an attracting Fatou coordinate defined on some petal P. It is easy to verify, using the injectivity of ϕ_A on P, that two points z_1 and z_2 are equivalent if and only if $\phi_A(z_1) - \phi_A(z_2) \in \mathbb{Z}$. Hence, ϕ_A defines by passage to quotients an injective mapping $\tilde{\phi}_A$ from $\mathcal{C}^\nu_A$ to $\mathbb{C}/\mathbb{Z}$. If we take P to be an ample petal, then it follows from Proposition 2.7 that $\tilde{\phi}_A$ takes on all values in $\mathbb{C}/\mathbb{Z}$. Thus:

Proposition 2.8. *The map $\tilde{\phi}_A$ is a conformal isomorphism from the Riemann surface $\mathcal{C}^\nu_A$ to the $\mathbb{C}/\mathbb{Z}$.*

In light of the preceding proposition, we will call $\mathcal{C}^\nu_A$ the *attracting cylinder* corresponding to the direction ν.

The *repelling cylinder* $\mathcal{C}_R$ for f is the attracting cylinder for a local inverse of f; by remarks just made, different choices of local inverse give the same repelling cylinder.

If P is an attracting petal, we will call the half-open domain

$$C_A = P \setminus f(P)$$

an attracting *fundamental crescent*, the name reflecting its shape. From the requirement that f^n converge uniformly to 0 on P, it follows that no point of P admits an infinite backward orbit in P. Thus, every point of P lies on an orbit starting in C_A, and distinct points of C_A lie on different orbits, so C_A serves as a fundamental domain for the orbits which converge to 0 through P. A repelling fundamental crescent C_R is defined similarly.

We note the following standard fact:

Proposition 2.9. *Let $h : \mathbb{C}^* \to \mathbb{C}^*$ be an injective holomorphic map. Then either $h(z) = cz$ or $h(z) = c/z$ for a non-zero constant c.*

Proof. Such an h is in particular an analytic function with isolated singularities at 0 and ∞ (and nowhere else.) By injectivity, neither singularity can be essential, so h extends to a meromorphic mapping of the Riemann sphere to itself, i.e. to a rational function. Injectivity on the sphere with two points deleted implies that this rational function has degree one, i.e., is a Möbius transformation. In particular, the extended function maps the sphere bijectively to itself, so either $h(0) = 0$, in which case $h(\infty) = \infty$, or $h(0) = \infty$. In the first case, $h(z)/z$ is bounded at ∞ and has a removable singularity at 0, so, by Liouville's Theorem, $h(z)/z = c$. In the case $h(0) = \infty$, applying the above to $h(1/z)$ gives $h(z) = c/z$.

□

A corollary of the above result is a uniqueness statement for Fatou coordinates:

Proposition 2.10. *Let P be an attracting petal of f, and let ϕ_1 and ϕ_2 be attracting Fatou coordinates on P, i.e., univalent analytic functions satisfying $\phi_i(f(z)) = \phi_i(z) + 1$. Then $\phi_2(z) - \phi_1(z)$ is constant on P.*

Proof. By Proposition 2.8, ϕ_1 and ϕ_2 both induce conformal isomorphisms from $\mathcal{C}_A$ to $\mathbb{C}/\mathbb{Z}$. It will be more convenient to work with with the punctured plane $\mathbb{C}^*$ instead of $\mathbb{C}/\mathbb{Z}$. The function

$$\operatorname{ixp}(z) = \exp(2\pi i z)$$

induces – again by passage to quotients – a conformal isomorphism from $\mathbb{C}/\mathbb{Z}$ to C^*, so $\operatorname{ixp} \circ \phi_1$ and $\operatorname{ixp} \circ \phi_2$ both induce conformal isomorphism $\mathcal{C}_A \to \mathbb{C}^*$. Hence, the prescription

$$h : \exp(2\pi i \phi_1(z))) \mapsto \exp(2\pi i \phi_2(z)) \quad \text{for all } z \in P$$

defines a conformal isomorphism $h : \mathbb{C}^* \to \mathbb{C}^*$. By Proposition 2.9, there are two possibilities:

(1) there is a non-zero constant – which we write as $\exp(2\pi i c)$ – so that

$$\exp(2\pi i c) \cdot \exp(2\pi i \phi_1(z) = \exp(2\pi i \phi_2(z)) \quad \text{for all } z \in P$$

or

(2) there is a constant c so that

$$\exp(2\pi i c)\cdot\exp(-2\pi i\phi_1(z) = \exp(2\pi i\phi_2(z)) \quad \text{for all } z\in P$$

In the first case,

$$\phi_2(z)-\phi_1(z)-c\in\mathbb{Z} \quad \text{for all } z\in P.$$

But the expression on the left is continuous, and an integer-valued continuous function on a connected set must be constant, so

$$\phi_2(z)=\phi_1(z)+c+n_0 \quad \text{for all } z\in P, \text{ for some } n_0\in\mathbb{Z}$$

which is what we wanted to prove.

In the second case, similarly,

$$\phi_2(z)=-\phi_1(z)+c+n_0 \quad \text{for all } z\in P,$$

and this contradicts

$$\phi_1(f(z))-\phi_1(z)=1=\phi_2(f(z))-\phi_2(z) \quad \text{for } z\in P,$$

so this case is excluded. □

For completeness, let us note how the situation changes if $f'(0)$ is a q-th root of unity $e^{2\pi ip/q}$ with $q\neq 1$. A fixed petal for the iterate f^q corresponds to a cycle of q petals for f. It thus follows that q divides the number n of attracting/repelling directions of 0 as a fixed point of f^q.

2.2. Asymptotic expansion of a Fatou coordinate at infinity.

We will now specialize to the case $q=1$ and $n=1$. By rescaling we can then bring the coefficient of z^2 to 1, and the normal form (2.2) becomes

$$f(z)=z+z^2+\alpha z^3+O(z^4). \tag{2.5}$$

We will say in this case that 0 is a *simple* parabolic fixed point of f. There is one attracting direction (-1) and one repelling direction $(+1)$. The change of variables κ moving the parabolic point to ∞ becomes simply

$$\kappa(z)=-\frac{1}{z},\quad \kappa^{-1}(w)=-\frac{1}{w},$$

and we have

$$F(w)=-\left(f(-\frac{1}{w})\right)^{-1}=w+1+\frac{A}{w}+O(w^{-2}),\quad \text{with } A=1-\alpha,$$

and $F(w)-w$ is analytic at ∞.

We showed earlier (Proposition 2.5) that any attracting Fatou coordinate at infinity Φ_A for such an f satisfies

$$\Phi_A(w) = w + o(w) \quad \text{as } w \to \infty \text{ appropriately}$$

We will prove shortly a much more precise result – an asymptotic expansion giving Φ_A up to corrections of order w^{-n} for any n. Before we do this, we investigate formal solutions to the functional equation

$$\Phi(F(w)) = \Phi(w) + 1$$

satisfied by both attracting and repelling Fatou coordinates.

Proposition 2.11. *There is a unique sequence $b_1, b_2, \ldots$ of complex coefficients such that*

$$\Phi_{ps}(w) = w - A\log(w) + \sum_{j=1}^{\infty} b_j w^{-j} \tag{2.6}$$

satisfies

$$\Phi_{ps} \circ F(w) = \Phi_{ps}(w) + 1 \quad \textit{in the sense of formal power series.} \tag{2.7}$$

Furthermore, if we set

$$\Phi_n(w) := w - A\log w + \sum_{j+1}^{n} b_j w^{-j} \tag{2.8}$$

then

$$\Phi_n(F(w)) - \Phi_n(w) - 1 = O(w^{-(n+2)}) \tag{2.9}$$

The logarithm appearing in (2.6) – and all other logarithms in this section – are to be understood as the *principal branch,* i.e., the branch with a cut along the negative real axis and real values on the positive axis. Because of the logarithmic term, Φ_{ps} as written is not exactly a formal power series in w^{-1}. To work around this, we rewrite the equation $\Phi_{\mathrm{ps}} \circ F = \Phi_{\mathrm{ps}} + 1$ formally as

$$F(w) - w - 1 - A\log(F(w)/w) = \sum_{j=1}^{\infty} b_j \left(F(w)^{-j} - w^{-j}\right) \tag{2.10}$$

Since $F(w)/w$ is analytic at ∞ and takes the value 1 there, $\log(F(w)/w)$ is analytic at ∞ and vanishes there. Furthermore, the formal identity

$$\log(F(w)) - \log(w) = \log(F(w)/w)$$

holds literally on $\{-\alpha < \operatorname{Arg}(w) < \alpha, |w| > R\}$ for sufficiently large R, for any $\alpha < \pi$. It is equation (2.10) which we really solve.

The left-hand side of (2.10) is analytic at ∞, and vanishes to second order there:

$$F(w)-w-1 = -Aw^{-1}+O(w^{-2}) \quad \text{and} \quad \log(F(w)/w) = w^{-1}+O(w^{-2})$$

Further, $F(w)^{-j}-w^{-j}$ is analytic at infinity and vanishes to order $j+1$ there, so

$$\sum_{j=1}^{\infty} b_j \left(F(w)^{-j} - w^{-j}\right)$$

is indeed a formal power series in w^{-1} which begins with a term in w^{-2}. Furthermore, the coefficient of w^{-j-1} in the expression on the right in (2.10) can be written as

$$-jb_j + \text{a function of } b_1,\, \ldots,\, b_{j-1}$$

Thus, since the left-hand side of (2.10) is known, the b_j's can be determined successively, and, by induction on j, they are uniquely determined. The assertion about the order of the error term $\Phi_n(F(w)) - \Phi_n(w) - 1$ also follows, since

$$\sum_{j=n+1}^{\infty} b_j \left(F(w)^{-j} - w^{-j}\right)$$

begins with a term in $w^{-(n+2)}$

Theorem 2.12. *Let b_1, b_2, … be as in Proposition 2.11, and let Φ_A be an attracting Fatou coordinate at infinity. Then, for any n,*

$$\Phi_A(w) = w - A\log w + C_A + \sum_{j=1}^{n} b_j w^{-j} + O(|w|^{-(n+1)}) \tag{2.11}$$

uniformly as $w \to \infty$ in any sector $-\alpha < \operatorname{Arg}(w) < \alpha$ with $\alpha < \pi$

We collect the main estimates needed for the proof of Theorem 2.12 in the following lemma:

Lemma 2.13.

$$\sum_{j=0}^{\infty} |F^j(w)|^{-m} = O(|w|^{-(m-1)}) \tag{I}$$

and

$$\left|(F^j)'(w)\right| \quad \text{is bounded uniformly in } j \tag{II}$$

both estimates holding uniformly for $w \to \infty$ in any sector $\{-\alpha < \operatorname{Arg}(w) < \alpha\}$ with $\alpha < \pi$.

Proof. We fix an $\alpha < \pi$, and we choose an α_1 with $\alpha < \alpha_1 < \pi$; it saves trouble later if we also require that $\pi - \alpha_1 < \pi/6$. Next we fix an R_0 large enough so that

$$|F(w) - (w+1)| < \sin(\pi - \alpha_1) \tag{2.12}$$

holds for $|w| > R_0$; then an R_1 so that the translated sector

$$\Delta(\alpha_1, R_1) := \{w : -\alpha_1 < \operatorname{Arg}(w - R_1) < \alpha_1\}$$

does not intersect the disk of radius R_0 about 0. Then (2.12) holds on $\Delta(\alpha_1, R_1)$, so F maps $\Delta(\alpha_1, R_1)$ to itself; also, since $\pi - \alpha_1 < \pi/6$, we also have – again from (2.12) –

$$|F(w) - w - 1| < \frac{1}{2} \quad \text{on } \Delta(\alpha_1, R_1). \tag{2.13}$$

from which it follows that

$$\operatorname{Re}(F(w) \geq \operatorname{Re}(w) + 1/2 \quad \text{for } w \in \Delta(\alpha_1, R_1). \tag{2.14}$$

and also

$$-(\pi - \alpha_1) < \operatorname{Arg}(F(w) - w) < (\pi - \alpha_1) \tag{2.15}$$

We will prove estimates (I) and (II) for $w \to \infty$ in $\Delta(\alpha_1, R_1) \cap \{-\alpha < \operatorname{Arg}(w) < \alpha\}$; this does what we want since every w with $-\alpha < \operatorname{Arg}(w) < \alpha$ and sufficiently large modulus is in $\Delta(\alpha_1, R_1)$.

Changing notation slightly: We want then to estimate

$$\sum_{j=0}^{\infty} |F^j(w_0)|^{-m},$$

for large w_0, given that

$$-\alpha_1 < \operatorname{Arg}(w_0 - R) < \alpha_1 \quad \text{and} \quad -\alpha < \operatorname{Arg}(w_0) < \alpha$$

where – crucially – $\alpha_1 > \alpha$. We write

$$F^j(w_0) =: w_j =: u_j + iv_j$$

In the calculation which follows, we adopt the convention that K denotes some constant depending only on α, α_1 and m. Different instances of K are not necessarily the same constant. All inequalities involving w_0 are only asserted to hold for $|w_0|$ large enough.

We first treat the case $-\pi/4 \leq \operatorname{Arg}(w_0) \leq \pi/4$, i.e., $|v_0| \leq u_0$. Then, by (2.14),

$$|w_j| \geq u_j \geq u_0 + j/2,$$

so

$$\sum_{j=0}^{\infty} |w_j|^{-m} \leq \sum_{j=0}^{\infty} (u_0 + j/2)^{-m} \leq K u_0^{-(m-1)} \leq K |w_0|^{-(m-1)};$$

in the last step, we used $|v_0| \leq u_0$ to estimate $u_0 \geq |w_0|/\sqrt{2}$. This proves the desired estimate in this case.

There remain the possibilities $\pi/4 < \mathrm{Arg}(w_0) < \alpha$ and $-\alpha < \mathrm{Arg}(w_0) < -\pi/4$. The estimates in the two cases are essentially the same; for definiteness we assume that the first holds, i.e., that w_0 is in the upper half-plane. By (2.15) the w_j are all contained in the translated sector $\{-(\pi-\alpha_1) < w - w_0 < +(\pi - \alpha_1)\}$. This sector intersects the diagonal line $\{\mathrm{Re}(w) = \mathrm{Im}(w)\}$ in a segment $(\hat{w}_-, \hat{w}_+)$ (labeled so that $|\hat{w}_-| < |\hat{w}_+|$.) It is easy to show, either by elementary geometry or by writing explicit formulas, that

$$|\hat{w}_-| \geq K^{-1}|w_0| \quad \text{and} \quad |\hat{w}_+| \leq K|w_0|.$$

Since $u_{j+1} \geq u_j + 1/2$, w_j, which start out above the diagonal line, will get and stay below it after finitely many steps. Let j_0 be the first index for which w_j is below the diagonal. Using the above upper bound on $|\hat{w}_+|$ we get a bound

$$j_0 \leq K|w_0|$$

The line segment from w_{j_0-1} to w_{j_0} intersects the diagonal line between $\hat{w}_-$ and $\hat{w}_+$. Since

$$|w_{j_0} - w_{j_0-1}| = |F(w_{j_0-1}) - w_{j_0-1}| < 3/2$$

we get

$$|w_{j_0}| \geq |\hat{w}_-| - 3/2 \geq K^{-1}|w_0| - 3/2 \geq K^{-1}|w_0|$$

By the first case treated

$$\sum_{j=j_0}^{\infty} |w_j|^{-m} \leq K|w_{j_0}|^{-(m-1)}| \leq K|w_0|^{-(m-1)} \tag{2.16}$$

All the w_j lie above (or on) the line through w_0 with direction $\alpha_1 - \pi$ (and the origin lies below this line.) Using $\mathrm{Arg}(w_0) < \alpha < \alpha_1$, the distance from this line to the origin satisfies a lower bound $K^{-1}|w_0|$. Hence each $|w_j| \geq K^{-1}|w_0|$. Since $j_0 \leq K|w_0|$,

$$\sum_{j=0}^{j_0-1} |w_j|^{-m} \leq j_0 \cdot \left(K^{-1}|w_0|\right)^{-m} \leq (K|w_0|) \cdot (K^m|w_0|^{-m}) \leq K|w_0|^{-(m-1)}$$

Combining this estimate on the sum of the first j_0 terms with the estimate (2.16) on the sum of the rest gives

$$\sum_{j=0}^{\infty} |F^j(w_0)|^{-m} \leq K|w_0|^{-(m-1)}$$

(for sufficiently large $|w_0|$), so (I) is established. (II) follows easily from (I) together with the estimate

$$F'(w) = 1 + O(|w|^{-2}),$$

the chain rule, and standard manipulations for reducing estimates on products to estimates on sums.

□

Proof of Theorem 2.12. Let $\pi/2 < \alpha < \pi$. By an argument already used several times, we can choose R sufficiently large so that

$$|F(w) - w - 1| < \sin(\pi - \alpha) \text{ and } \operatorname{Re}(F(w) > \operatorname{Re} w + 1/2$$

for all $w \in \Delta(\alpha, R) := \{-\alpha < \operatorname{Arg}(w - R) < \alpha\}$ As usual, it follows from the first of these inequalities that $F(\overline{\Delta(\alpha, R)}) \subset \Delta(\alpha, R)$. We are going to prove

$$\Phi_A(w) = \Phi_n(w) + O(|w|^{-(n+1)}) \quad \text{as } w \to \infty \text{ in } \Delta(\alpha, R)$$

(where Φ_n is defined by (2.7)); this assertion for all α implies the assertion of the theorem for all α.

We set

$$c_n(w) := \Phi_A(w) - \Phi_n(w) \quad \text{and} \quad u_n(w) := \Phi_n \circ F(w) - \Phi_n(w) - 1.$$

By Proposition 2.11, u_n is analytic at infinity with

$$u_n(w) = d_{n+2} w^{-(n+2)} + \cdots . \tag{2.17}$$

A simple calculation gives

$$c_n \circ F(w) = c_n(w) - u_n(w);$$

iterating gives

$$c_n \circ F^k(w) = c_n(w) - \sum_{j=0}^{k-1} u_n(F^j(w));$$

reorganizing and differentiating gives

$$c_n'(w) = \sum_{j=0}^{k-1} u_n'(F^j(w)) \cdot (F^j)'(w) + c_n'(F^k(w)) \cdot (F^k)'(w). \tag{2.18}$$

By differentiating (2.17)

$$u_n'(w) = O(|w|^{-(n+3)}) \text{ as } w \to \infty,$$

so, by Lemma 2.13,

$$\sum_{j=0}^{\infty} \left| u_n'(F^j(w)) \cdot (F^j)'(w) \right| < \infty$$

By Proposition 2.5

$$\Phi_A'(w) \to 1 \quad \text{as } \operatorname{Re}(w) \to +\infty,$$

and the same is true for Φ_n by an elementary calculation; hence

$$c_n'(w) \to 0 \quad \text{as } \operatorname{Re}(w) \to +\infty.$$

Thus, we can let $k \to \infty$ in (2.18) to get

$$c_n'(w) = \sum_{j=0}^{\infty} u_n'(F^j(w)) \cdot (F^j)'(w)$$

Applying both parts of Lemma 2.13 to this representation,

$$|c_n'(w)| \leq \text{const}\, |w|^{-(n+2)} \tag{2.19}$$

for all $w \in \Delta(\alpha, R)$. It follows from this estimate that the limit

$$\lim_{u \to \infty} c_n(u + iv)$$

exists and is independent of v. We denote this limit by C_A. Then

$$c_n(u + iv) = C_A - \int_u^{\infty} c_n'(\sigma + iv) d\sigma,$$

so by integrating (2.19) we get

$$|c_n(w) - C_A| \leq \text{const}\, |w|^{-(n+1)}$$

for all $w \in \Delta(\alpha, R)$, which is what we set out to prove.

□

2.3. **Écalle-Voronin invariants and definition of parabolic renormalization.** The Riemann surface $\mathbb{C}/\mathbb{Z}$ has two punctures at the upper end ($\operatorname{Im} z \to +\infty$), and at the lower end ($\operatorname{Im} z \to -\infty$). Filling them with points $\oplus$ and $\ominus$ respectively, we obtain the Riemann sphere. The mapping

$$\operatorname{ixp}(z) \equiv \exp(2\pi i z),$$

conformally transforms $\mathbb{C}/\mathbb{Z} \mapsto \mathbb{C}^*$, sending $\oplus \to 0$ and $\ominus \to \infty$.

Consider f with a simple parabolic fixed point. The intersection $P_R \cap P_A$ projects to a pair of "polar caps": open regions $W^+ \ni \oplus$ and $W^- \ni \ominus$ in the repelling cylinder $\mathcal{C}_R$. Consider the pair of analytic maps

$$h : W^+ \cup W^- \to \mathcal{C}_A, \text{ given by } h \equiv \phi_A \circ \phi_R^{-1}.$$

The asymptotics in Lemma 2.13 implies that

$$\operatorname{Im} h(w) \to \pm\infty \text{ when } \operatorname{Im} w \to \pm\infty.$$

Thus both singularities $\oplus$, $\ominus$ are removable for h. It is easier to discuss this on the Riemann sphere, so let us denote

$$H \equiv \operatorname{ixp} \circ h \circ \operatorname{ixp}^{-1} = \operatorname{ixp} \circ \phi_A \circ \phi_R^{-1} \circ \operatorname{ixp}^{-1} .$$

This is an analytic mapping in $\operatorname{ixp}(W^+ \cup W^-)$ which has removable singularities at 0 and ∞. The germ of $H(z)$ at 0 and ∞ is not uniquely defined, of course, as there is some freedom in the definition of Fatou coordinates. By Proposition 2.10, $h(w)$ is defined up to a pre- and post-composition with translations. Thus, $H(w)$ may be replaced with $cH(dw)$ for some $c, d \in \mathbb{C}^*$. It is elementary to see that the product $H'(0) \cdot H'(\infty)$ is independent of the choice of the Fatou coordinates. In fact, a calculation (see [BE]) shows that:

Proposition 2.14. *The quantity*

$$H'(0) \cdot H'(\infty) = e^{4\pi^2 A}, \text{ where } A = 1 - \alpha, \text{ as before.}$$

Proof. Let us write the two-term asymptotic developments for the attracting and repelling Fatou coordinates respectively as:

$$g_A(w) = w - A \log_A(w) + \text{const}_A + O\left(\frac{1}{|w|}\right),$$

$$g_R(w) = w - A \log_R(w) + \text{const}_R + O\left(\frac{1}{|w|}\right),$$

where $\log_A$ and $\log_R$ denote the corresponding branches of the log. Since we are free to choose both the constants and the logs without changing the value of the product $H'(0) \cdot H'(\infty)$, let us do this as follows. We set $\text{const}_A = \text{const}_R = 0$. For $\log_R$ we select the branch of the log in $\mathbb{C} \setminus \mathbb{R}_+$ which takes the value $i\pi$ at -1; for $\log_A$ we select the branch defined in $\mathbb{C} \setminus \mathbb{R}_-$ with $\log_A(1) = 0$. Now,

$$\lim_{\operatorname{Im} w \to -\infty} g_A(w) - g_R(w) = -2\pi i A, \text{ and } \lim_{\operatorname{Im} w \to +\infty} g_A(w) - g_R(w) = 0.$$

Exponentiating, we get

$$H'(0) \cdot H'(\infty) = e^{4\pi^2 A}.$$

□

Let us use the notation $H_f(z)$ to denote the pair of analytic germs $H(z)$ at $z = 0$ and $z = \infty$ for a given parabolic germ $f(z) = z + z^2 + \cdots$. The following theorem is due to Voronin [Vor] (an equivalent version was proved by Écalle [Ec]):

Theorem 2.15. *Two analytic germs $f(z)$, $g(z)$ of the form (2.5) are conjugated by a local conformal change of coordinates $\varphi(z)$ with $\varphi(0) = 0$ if and only if*

$$H_f(z) \equiv cH_g(dz) \text{ for } c, d \in \mathbb{C}^*. \tag{2.20}$$

Sketch of proof. Denote the objects corresponding to one of the maps f, g by a corresponding superscript: $\mathcal{C}^f_A$, ϕ^f_A, etc. To prove the "only if" direction, consider a topological disk $U \ni 0$ in which the conformal conjugacy φ is defined. Let $P^f_A \cup P^f_R$ be a pair of attracting/repelling petals contained in U. Then $P^g_A = \varphi(P^f_A)$, $P^g_R = \varphi(P^f_R)$ form a Leau-Fatou flower for g. As a conjugacy, φ maps orbits of f to orbits of g, and hence induces conformal mappings

$$\hat{\varphi}_A : \mathcal{C}^f_A \to \mathcal{C}^g_A, \text{ and } \hat{\varphi}_R : \mathcal{C}^f_R \to \mathcal{C}^g_R.$$

By Liouville's Theorem, these mappings are translations. Under the conformal isomorphism $\text{ixp} : \mathbb{C}/\mathbb{Z} \to \hat{\mathbb{C}}$, they become complex multiplications

$$\tilde{\varphi}_A(\zeta) = a\zeta, \text{ and } \tilde{\varphi}_R(\zeta) = b\zeta.$$

Since $\varphi \circ f \circ \varphi^{-1} = g$, we have

$$H_g(z) = \tilde{\varphi}_A \circ H_f \circ \tilde{\varphi}_R(z) = aH_f(z/b).$$

Conversely, starting with the equation (2.20) let us normalize the Fatou coordinates for f and g so that

$$H_f \equiv H_g. \tag{2.21}$$

We can define a conjugacy φ between f and g by setting it equal to $(\phi^g_A)^{-1} \circ \phi^f_A$ in P^f_A and to $(\phi^g_R)^{-1} \circ \phi^f_R$ in P^f_R. The equation (2.21) the two definitions coincide in the intersection $P^f_A \cap P^f_R$. □

Therefore, the conformal conjugacy classes of parabolic germs of the form (2.5) are completely determined by the pair of analytic germs $\mathcal{E}_f$. The latter can, in fact, be selected arbitrarily:

Theorem 2.16. *Let $H(z)$ be any pair of analytic germs defined at 0 and at ∞, with $H'(0)H'(\infty) \neq 0$. Then there exists a simple parabolic germ $f(z)$ of the form*

$$f(z) = z + z^2 + \alpha z^3 + \cdots ,$$

for which

$$H_f \equiv H.$$

Moreover,

$$H'(0)H'(\infty) = e^{4\pi^2(1-\alpha)}.$$

We leave out the proof; it can be obtained, for instance, using a quasiconformal surgery construction and Measurable Riemann Mapping Theorem.
We will now make several key definitions.

Definition 2.2. We say that a parabolic germ $f(z)$ of the form (2.2) is *renormalizable* if there exists a (necessarily unique)choice of non-zero constants c and d such that

$$\mathcal{E}_f \equiv cH_f(dz) = z + z^2 + \cdots$$

near the origin.

That is, for a renormalizable map f, there is a choice of normalization of the Fatou coordinates, which makes the point 0 for $H_f(z)$ a *simple* parabolic fixed point.

Definition 2.3. For a renormalizable germ $f(z)$ as in (2.5) we will call the analytic germ of $\mathcal{E}_f$ at $z = 0$ the *parabolic renormalization* of f. We will use the notation $\mathcal{P}(f)$ for the parabolic renormalization.

For the appropriately normalized choice of ϕ_A and ϕ_R we thus have:

$$\mathcal{P}(f) = \operatorname{ixp} \circ \phi_A \circ (\phi_R)^{-1} \circ \operatorname{ixp}^{-1}, \tag{2.22}$$

with suitably selected branches of the inverses.

3. Global theory

3.1. **Basic facts about branched coverings.** We give a brief summary of relevant facts about analytic branched coverings. A more detailed exposition is found e.g. in the Appendix E of [Mil1].

For a holomorphic map $f : X \to Y$ between two Riemann surfaces a *regular value* is a point $y \in Y$ for which one can find an open neighborhood $U = U(y)$ such that

$$f^{-1}(U) \xrightarrow{f} U$$

is a covering map. The complement of this set consists of the *singular values* of f, and will be denoted $\operatorname{Sing}(f)$.

By definition, $y \in Y$ is an *asymptotic value* of f if there exists a parametrized path

$$\gamma : (0, 1) \to X, \text{ such that } \lim_{t\to 1-} f(\gamma(t)) = y,$$

and such that $\lim_{t\to 1-} \gamma(t)$ does not exist in X. We will be concerned with the situation when X is a proper subdomain of the Riemann sphere. In this case, the non-existence of the limit can be replaced with:

$$\gamma(t) \underset{t\to 1-}{\longrightarrow} \partial X.$$

We will denote $\operatorname{Asym}(f) \subset Y$ the set of all asymptotic values of f.

Recall that $y_0 \in Y$ is called a *critical value* (or a *ramified point*) of f if there exists $x_0 \in X$ such that $y_0 = f(x_0)$, and the local degree of f at x_0 is $n \geq 2$. Thus, in local coordinates, one has

$$f(x) - y_0 = c(x - x_0)^n + \cdots \text{ where } n \geq 2.$$

The point x_0 is called a *critical point* of f (or a *ramification point*). We denote $\operatorname{Crit}(f) \subset X$ the set of the critical points of f. One then has:

Proposition 3.1. *For a holomorphic map $f : X \to Y$ between Riemann surfaces,*

$$\operatorname{Sing}(f) = \overline{\operatorname{Asym}(f) \cup f(\operatorname{Crit}(f))};$$

that is, the set of the singular values of f is the closure of the union of its asymptotic and critical values.

Recall that a non-constant map $g : S_1 \to S_2$ between two Hausdorff topological spaces is called *proper* if the preimage of every compact set of S_2 is compact in S_1. It is easy to see that:

Proposition 3.2. *Let $f : X \to Y$ be a proper holomorphic map of Riemann surfaces. Let W be a non-empty connnected open subset of Y, and let V be a connected component of $f^{-1}(W)$. Then $f : V \to W$ is also proper.*

For a holomorphic map $f : X \to Y$ define the degree of f at $y \in Y$ (denoted $\deg_y(f)$) as the possibly infinite sum of the number of preimages of y in X, counted with multiplicity. It is not difficult to see that a proper analytic map has a well-defined local degree:

Proposition 3.3. *If $f : X \to Y$ is a proper analytic map between Riemann surfaces, and Y is connected, then $\deg_y(f)$ is finite and independent of y (and can be denoted as $\deg(f)$).*

As a consequence, note:

Proposition 3.4. *If $f : X \to Y$ is a proper analytic map between Riemann surfaces, and Y is connected, then $f(X) = Y$.*

Proposition 3.5. *Let $g : S_1 \to S_2$ be a proper continuous map between Hausdorff topological spaces which is everywhere a local homeomorphism. Then f is a covering map.*

In particular, putting together Proposition 3.3 and Proposition 3.5, we have

Proposition 3.6. *Let $f : X \to Y$ be a proper analytic mapping between Riemann surfaces. Let Y be connected, and set $d = \deg(f)$. Then*

$$f : X \setminus \mathrm{Crit}(f) \to Y \setminus f(\mathrm{Crit}(f))$$

is a degree d covering.

In view of the above, a proper analytic map is sometimes called a *branched covering of a finite degree.* Generalizing to the case when local degree is infinite gives the following definition:

Definition 3.1. A holomorphic map $f : X \to Y$ between Riemann surfaces is a *branched covering* if every point $y \in Y$ has a connected neighborhood $U \equiv U(y)$ such that the restriction of f to each connected component of $f^{-1}(U)$ is a proper map.

As a canonical example, consider the case $X = Y = \hat{\mathbb{C}}$. Non-constant rational maps f are clearly branched coverings; and every branched covering is, in fact, a rational map.

Let us formulate another general lemma which we will find useful:

Lemma 3.7. *Let $f : U \to V$ be a proper analytic map between connected subdomains of $\hat{\mathbb{C}}$. Assume that f has a single critical value $v \in V$, and that V is a topological disk. Then U is also a topological disk, and f has only one critical point u in U, such that $f^{-1}(v) = \{u\}$.*

Proof. Set $\hat{V} \equiv V \setminus \{v\}$ and $\hat{U} \equiv f^{-1}(\hat{V})$. By Proposition 3.6, the map $f : \hat{U} \to \hat{V}$ is a covering. The domain $\hat{V}$ is homeomorphic to $\mathbb{D} \setminus \{0\}$. By the standard facts about coverings of $\mathbb{D} \setminus \{0\}$, the domain $\hat{U}$ is also homeomorphic to the punctured disk. Moreover, denoting $\eta_{\hat{V}}$ and $\eta_{\hat{U}}$ conformal homeomorphisms mapping $\mathbb{D} \setminus \{0\}$ to $\hat{V}$ and $\hat{U}$ respectively, we see that

$$\eta_{\hat{V}}(f(z)) = \mathrm{const} \cdot (\eta_{\hat{U}}(z))^d \text{ for } d \geq 2.$$

□

3.2. **Parabolic renormalization of the quadratic map.** As a key example, consider the quadratic polynomial

$$f_0(z) = z \mapsto z + z^2.$$

By the Fatou-Sullivan Classification, the filled Julia set $K(f_0)$ coincides with the closure of the basin of 0:

$$K(f_0) = \overline{\{z |\ f^n(z) \to 0\}}.$$

Through the functional equation

$$\phi_A(f_0(z)) = \phi_A(z) + 1,$$

the attracting Fatou coordinate extends to a map

$$\phi_A : K(f_0) \to \mathbb{C}.$$

It is straightforward to show that (cf. [Do2]):

Proposition 3.8. *The map $\phi_A : K(f_0) \to \mathbb{C}$ is a branched covering. The critical points of ϕ_A are the preimages of the critical value $-1/4$ of f_0:*

$$\mathrm{Crit}(\phi_A) = \cup_{n\geq 1} f_0^{-n}(-1/4).$$

All of them have local degree 2, and their images are integer translates of each other:

$$\phi_A(\mathit{Crit}(\phi_A)) = \{\phi_A(-1/4) + \mathbb{Z}\}.$$

Note that the closure

$$\overline{\mathrm{Crit}(\phi_A)} = \mathrm{Crit}(\phi_A) \cup J(f_0). \tag{3.1}$$

We prove the following:

Theorem 3.9. *The germs H_{f_0} at 0 and ∞ have maximal analytic continuations to the two Jordan domains $\hat{W}^+ \subset \hat{\mathbb{C}}$ and $\hat{W}^- \subset \hat{\mathbb{C}}$ respectively. These domains are the two connected components of the set*

$$\hat{W} \equiv \mathrm{ixp} \circ \phi_R(K(f_0) \cap P_R) \cup \{0, \infty\}.$$

Definition 3.2. Let $g : U \to \hat{\mathbb{C}}$ be an analytic map of an open (not necessarily connected) domain U. Suppose g has a maximal analytic continuation to a domain $W \supset U$. We will write $W = \mathrm{Dom}(g)$.

Thus $\mathrm{Dom}(H_{f_0}) = \hat{W}^+ \cup \hat{W}^-$ (see Figure 5.3).

Theorem 3.9 relies on a well-known folklore result:

Theorem 3.10. *The Julia set of the quadratic polynomial $f_0(z) = z + z^2$ is a Jordan curve. The dynamics of f_0 restricted to the Julia set is topologically conjugated to the angle-doubling map of the circle. Specifically, there exists a unique continuous map $\rho : J(f_0) \to \mathbb{T}$ such that*

$$\rho(f_0(z)) = 2\rho(z) \bmod 1.$$

In the next subsection we will provide a proof of Theorem 3.10; while it is fairly standard, the strategy used in it will be useful to us in what follows. Let us proceed with proving Theorem 3.9, assuming Theorem 3.10:

Proof of Theorem 3.9. Standard considerations of Montel's Theorem imply that the topological disk $B_0 \equiv \overset{\circ}{K}(f_0)$ is the basin of the parabolic fixed point 0. Thus, an orbit of a point $z_0 \in B_0$ eventually enters an attracting petal P_A. On the other hand, the orbit of a point $z_0 \in \hat{\mathbb{C}} \setminus \overline{B_0}$ converges to ∞, and hence

$$\mathrm{Dom}(H_{f_0}) = \mathrm{ixp} \circ \phi_R(K(f_0) \cap P_R) \cup \{0, \infty\} = \hat{W}.$$

Consider the unique real-analytic conformal map ψ which sends B_0 to the slit domain

$$W \equiv \hat{\mathbb{C}} \setminus (\mathbb{R}_{\geq 0} \cup \{\infty\}),$$

so that

$$\psi(-1/2) = -1 \text{ and } \lim_{x \to 0- \text{ and } x \in \mathbb{R}} \psi(x) = 0.$$

By Carathéodory Theorem, ϕ has a continuous extension to the boundary:

$$J(f_0) \to \mathbb{R}_{\geq 0} \cup \{\infty\} \subset \hat{\mathbb{C}}.$$

By real symmetry, the map

$$h \equiv \psi \circ f_0 \circ \psi^{-1} : W \to W$$

extends continuously, and hence, holomorphically, to $\hat{\mathbb{C}}$. The quadratic rational map h has a parabolic fixed point at the origin: this follows from the fact that on the real axis, 0 is attracting for h in the negative direction, and repelling in the positive one. It is not difficult to verify that

$$h(z) = \frac{z}{(z-1)^2}. \tag{3.2}$$

The conformal conjugacy ψ induces conformal isomorphisms between spaces of orbits:

$$\mathcal{C}_A^{f_0} \to \mathcal{C}_A^h, \text{ and}$$

$$\mathrm{Dom}(H_{f_0}) = \hat{W} \to \mathrm{Dom}(H_h) = \mathrm{ixp} \circ \phi_R^h(P_R^h \cap W) \cup \{0, \infty\}.$$

The Julia set

$$J(h) = \mathbb{R}_{\geq 0}.$$

By Montel's Theorem, every open set $U \cap \mathbb{R}_{\geq 0} \neq \emptyset$ contains preimages of all points in $\hat{\mathbb{C}}$, except at most two. Hence, H_h cannot be analytically extended across the "equator"

$$\mathrm{ixp} \circ \phi_R^h(\mathbb{R}_{>0}).$$

Thus, by considerations of real symmetry, $\mathrm{Dom}(H_h)$ is a union of disks around 0 and ∞.

This implies that the domain $\hat{W}$ is a union of two topological disks $\hat{W}^+$ and $\hat{W}^-$, whose boundaries lie in the projection $\mathrm{ixp} \circ \phi_R^{f_0}(J(f_0))$.

Suppose that one of the boundaries is not a Jordan curve; to fix the ideas, assume that it is $L \equiv \partial \hat{W}^+$. In view of Theorem 3.10, L is locally connected; in view of the above, it is also connected. In view of Carathéodory theory, there is a point $w \in L$, which is bi-accessible from $\hat{W}^+$. Let $z \in J(f_0)$ be a point with $\operatorname{ixp} \circ \phi_R^{f_0}(z) = w$. The map $\operatorname{ixp} \circ \phi_R^{f_0}$ is a local homeomorphism at z, hence the point z is bi-accessible from the parabolic basin of f_0. This contradicts Theorem 3.10, and the proof is thus completed. $\square$

We now proceed with the proof of Theorem 3.10

3.3. **The Julia set of $f_0(z) = z + z^2$ is a Jordan curve.** We begin by defining several useful local inverse branches of f_0, by choosing the appropriate branches of the square root in the formal expression

$$f_0^{-1}(w) = \sqrt{w + 1/4} - 1/2. \tag{3.3}$$

We first cut the plane along the positive real ray $R \equiv [-1/4, +\infty)$ and denote

$$g_0 : \mathbb{C} \setminus R \to \mathbb{H}$$

the inverse branch with non-negative imaginary part. Similarly, we define

$$g_1 : \mathbb{C} \setminus R \to -\mathbb{H}$$

to be the inverse branch with negative imaginary part, so that $g_1 \equiv -g_0$.

Finally, we define an inverse branch g which fixes the parabolic point $z = 0$. We slit the plane along the ray $(-\infty, -1/4]$, and select the branch of the square root in (3.3) with non-negative real part. In this way, we get

$$g : \mathbb{C} \setminus (-\infty, -1/4] \to \{\operatorname{Re} z \geq -1/2\}.$$

We note that Taylor expansion of g at $w = 0$ begins with $w - w^2 + \cdots$, and thus it also has a parabolic fixed point at the origin.

When needed, we will continuosly extend the three iverse branches defined above to the banks of the slits.

We note, that

Proposition 3.11. *The branch g maps $\mathbb{C} \setminus (-\infty, 0]$ into itself.*

Applying Denjoy-Wolf Theorem, we obtain

Corollary 3.12. *The successive iterates $(g)^n$ converge uniformly to the parabolic point 0 on compact subsets of $\mathbb{C} \setminus (-\infty, 0]$.*

Let us note:

Proposition 3.13. *The intersection*

$$J(f_0) \cap \mathbb{R} = \{-1, 0\}.$$

Proof. Indeed, if $x > 0$ then

$$f(x) = x + x^2 > x, \text{ and, furthermore, } f^n(x) > x + nx^2 \to \infty,$$

and hence $(0, \infty) \subset F(f_0)$. Since

$$f_0((-\infty, -1)) = (0, \infty),$$

the same holds for $(-\infty, -1)$. Finally, f_0 maps the interval $(-1, 0)$ to itself and $f_0(x) > x$ for x in this interval, so $f_0^n(x) \to 0$. Hence, $(-1, 0) \subset F(f)$ as well, as claimed. □

Define

$$\mathbf{A} := \{z : 1/2 < |z + 1/2| < 2\}.$$

We invite the reader to verify:

Proposition 3.14.

1. *if $f(z) \in \mathbf{A}$, then $z \in \mathbf{A}$, i.e., $f_0^{-1}\mathbf{A} \subset \mathbf{A}$.*
2. *$J(f_0)$ is contained in the closure $\overline{\mathbf{A}}$ of $\mathbf{A}$.*

To prove Theorem 3.10, we are going to show that $f_0^{-n}\mathbf{A}$ are a decreasing sequence of topological annuli which shrink down to a Jordan curve J. Points inside J are attracted to the parabolic point; those outside are attracted to ∞. Thus, the Jordan curve J contains the Julia set. The restriction of f_0 to J is topologically conjugate to $\theta \mapsto 2 \cdot \theta \bmod 1$ on the circle $\mathbb{R}/\mathbb{Z}$. In particular, periodic points are dense in J, so the Julia set is the whole of J.

Set

$$\mathbf{A}_0 := \{z \in \mathbf{A} \,:\, \operatorname{Im} z > 0\} \quad \text{and} \quad \mathbf{A}_1 := \{z \in \mathbf{A} \,:\, \operatorname{Im} z < 0\}$$

If $i_0 i_1 \ldots i_n$ is an arbitrary sequence of $n + 1$ 0's and 1's, we define

$$\mathbf{A}_{i_0 i_1 \ldots i_n} := \{z \in \mathbf{A}_0 : f^j(z) \in \mathbf{A}_{i_j} \quad \text{for} \quad 1 \le j \le n\}$$

It follows from the definitions that

- $\mathbf{A}_{i_0 i_1 \ldots i_n}$ is decreasing in n: $\mathbf{A}_{i_0 \ldots i_n} \subset \mathbf{A}_{i_0 \ldots i_{n-1}}$;
- $f\mathbf{A}_{i_0 i_1 \ldots i_n} = \mathbf{A}_{i_1 \ldots i_n}$;
- $\mathbf{A}_{i_0 i_1 \ldots i_n} = g_{i_0}\mathbf{A}_{i_1 \ldots i_n}$, and hence

  $$\mathbf{A}_{i_0 i_1 \ldots i_n} = g_{i_0} \circ g_{i_1} \circ \cdots \circ g_{i_{n-1}}\mathbf{A}_{i_n},$$

  or, more generally,

  $$\mathbf{A}_{i_0 i_1 \ldots i_n} = g_{i_0} \circ g_{i_1} \circ \cdots \circ g_{i_{j-1}}\mathbf{A}_{i_j i_{j+1} \ldots i_n} \quad \text{for } 1 \le j \le n - 1.$$

The main step in the argument is to show that the diameters of the puzzle pieces $\mathbf{A}_{i_0 i_1 \ldots i_n}$ go to zero as $n \to \infty$, uniformly in $i_0 i_1 \ldots i_n$. More precisely, let us define

$$\rho_n := \sup \left\{ \operatorname{diam}(\mathbf{A}_{i_0 \ldots i_n}) : i_0 \ldots i_n \in \{0,1\}^{n+1} \right\}$$

Since $\mathbf{A}_{i_0 \ldots i_n} \supset \mathbf{A}_{i_0 \ldots i_{n+1}}$, the sequence ρ_n is non-increasing in n, so

$$\rho_* := \lim_{n \to \infty} \rho_n \tag{3.4}$$

exists.

We will prove the following:

Proposition 3.15. *The limit* $\rho_* = 0$.

Before proving the Proposition, let us formulate a helpful Lemma:

Lemma 3.16. *There is an infinite sequence* $i_0 i_1 \ldots i_n \ldots$ *so that*

$$\operatorname{diam}(A_{i_0 i_1 \ldots i_n}) \geq \rho_* \quad \textit{for all } n,$$

i.e., a descending chain *of puzzle pieces whose diameters do not go to zero.*

Proof of Lemma 3.16. A simple diagonal argument implies the existence of a sequence $i_0 i_1 \ldots i_k \ldots$ such that, for all k,

$$\lim_{n \to \infty} \sup \left\{ \operatorname{diam}(\mathbf{A}_{i_0, \ldots i_k i_{k+1}} \ldots i_{k+n}) : i_{k+1} \ldots i_{k+n} \in \{0,1\}^n \right\} \geq \rho_*$$

Using the fact that $\operatorname{diam}(\mathbf{A}_{i_0 i_1 \ldots i_n})$ is non-increasing in n, we get

$$\operatorname{diam}(\mathbf{A}_{i_0 i_1 \ldots i_k}) \geq \rho_* \quad \text{for all } k,$$

proving the lemma. ☐

Proof of Proposition 3.15. Let us assume the contrary: $\rho_* > 0$.

We fix a sequence $i_0 i_1 \ldots$ as in the Lemma 3.16, and we split the proof into three cases:

(1) i_j is eventually 0;
(2) i_j is eventually 1;
(3) neither of the above holds.

We start with case (3). There are then infinitely many j's so that

$$i_j = 0 \quad \text{and} \quad i_{j+1} = 1$$

Let j_k be a strictly increasing sequence of such j's. Then, for each k,

$$f_{-j_k} := g_{i_0} \circ g_{i_1} \circ \cdots \circ g_{i_{j_k - 1}}$$

is an analytic branch of the inverse of $f_0^{j_k}$ mapping $\mathbf{A}_{01}$ bijectively to $\mathbf{A}_{i_0 \ldots i_{j_k + 1}}$. The closure of $\mathbf{A}_{01}$ does not intersect the postcritical set of f_0. The following argument is standard (see [Lyu], Proposition 1.10).

We let U be an simply connected open neighborhood of $\overline{\mathbf{A}_{01}}$ disjoint from the postcritical set. Then each f_{-j_k}'s extends to an analytic branch of the inverse of $f_0^{j_k}$ defined on U (and we denote this extension also by f_{-j_k}).

By Montel's Theorem, there is a subsequence of (j_k) along which f_{-j_k} converges uniformly on compact subsets of U in the spherical metric on $\hat{\mathbb{C}}$. By adjusting the notation, we can assume that the sequence (f_{-j_k}) itself converges to an analytic function which we denote h. Since

$$\operatorname{diam}(f_{-j_k}\mathbf{A}_{01}) = \operatorname{diam}(\mathbf{A}_{i_0 \dots i_{j_k+1}})$$

does not go to zero as $k \to \infty$, the function h is non-constant.

Let $z_0 := g_0(g_1(-1)) \in \mathbf{A}_{01} \cap J(f_0)$. By invarince of the Julia set, we have $w_0 = h(z_0) \in J(f_0)$. Since h is non-constant, $h(\mathbf{A}_{01})$ contains an open neighborhood W of w_0. Let $V \Subset W$ be a smaller open neighborhood of w_0. Then, an arbitrary large iterate $f_0^{j_k}$ maps V inside U, which contradicts Montel's theorem.

We turn next to case 1 above, and deal first with the situation $i_j = 0$ for all j. We write temporarily

$$\mathbf{A}^{(n)} := \mathbf{A}_{\underbrace{0 \cdots 0}_{n \text{ terms}}} = (g_0)^{n-2}\mathbf{A}_{00}$$

Now g_0 maps $\mathbf{A}_{00}$ into itself, and $g_0 = g$ on $\mathbf{A}_{00}$, so we can write

$$\mathbf{A}^{(n)} = (g)^{n-2}\mathbf{A}_{00}.$$

By Corollary 3.12 and local properties of parabolic dynamics near 0,

$$(g)^{n-2} \to 0 \quad \text{uniformly on } \mathbf{A}_{00},$$

so

$$\operatorname{diam}(\mathbf{A}^{(n)}) \to 0 \quad \text{as } n \to \infty.$$

Next consider sequences of the form $i_0 i_1 \cdots i_k 00 \cdots$, and use the formula

$$\mathbf{A}_{i_0 \dots i_k \underbrace{0 \cdots 0}_{n \text{ terms}}} = g_{i_0} \circ g_{i_1} \circ \cdots \circ g_{i_k} A^{(n)}.$$

The mapping $g_{i_0} \circ g_{i_1} \circ \cdots \circ g_{i_k}$ extends to be continuous on $\overline{\mathbf{A}_{00}}$, and $\operatorname{diam}(A^{(n)}) \to 0$ by what we just proved, so

$$\operatorname{diam}(\mathbf{A}_{i_0 \dots i_k \underbrace{0 \cdots 0}_{n \text{ terms}}}) \to 0 \quad \text{as } n \to \infty.$$

A similar argument, using $g_1 = g$ on $\mathbf{A}_{11}$ shows that

$$\operatorname{diam}(\mathbf{A}_{i_0 \dots i_k \underbrace{1 \cdots 1}_{n \text{ terms}}}) \to 0 \quad \text{as } n \to \infty.$$

□

Let $\mathbf{i} = (i_j)_{j=0}^{N}$ be a finite or infinite sequence ($N \leq \infty$) of 0's and 1's. We interpret it as a binary representation of a number in $[0, 1]$:

$$\underline{\mathbf{i}}_2 \equiv \sum_{j=0}^{N} i_j 2^j \in [0, 1].$$

For any such dyadic sequence,

$$\overline{\mathbf{A}_{i_0}} \supset \overline{\mathbf{A}_{i_0 i_1}} \supset \cdots \supset \overline{\mathbf{A}_{i_0 \ldots i_n}} \supset \ldots$$

is a nested sequence of compact sets in $\mathbb{C}$ with diameter going to 0, so its intersection contains exactly one point, which we denote by $\hat{z}(\mathbf{i})$. It is immediate from the construction that $\mathbf{i} \mapsto \hat{z}(\mathbf{i})$ is continuous from $\{0, 1\}^{\mathbb{Z}}$ to $\mathbb{C}$. It is not injective, however, this ambiguity is easily tractable:

Lemma 3.17. *Suppose* $\mathbf{i} = i_0 i_1 \ldots i_{n-1}$ *and* $\mathbf{i}' = i'_0 i'_1 \ldots i'_{n-1}$ *are two finite dyadic sequences of an equal length, and*

$$\overline{\mathbf{A}_{\mathbf{i}}} \cap \overline{\mathbf{A}_{\mathbf{i}'}} \neq \emptyset.$$

Then either $\underline{\mathbf{i}}_2 = \underline{\mathbf{i}'}_2$, *or* $\underline{\mathbf{i}}_2 = \underline{\mathbf{i}'}_2 \pm 2^{-n}$.

The proof is a straightforward induction in n, and is left to the reader.

As a corollary, we get:

Corollary 3.18. *Let* $\mathbf{i}$ *and* $\mathbf{i}'$ *be two infinite dyadic sequences. If If* $\underline{\mathbf{i}}_2 \neq \underline{\mathbf{i}'}_2$, *then* $\overline{\mathbf{A}_{i_0 \cdots i_n}}$ *and* $\overline{\mathbf{A}_{i'_0 \cdots i'_n}}$ *are disjoint for large enough* n

Proof. For any $\mathbf{i} = i_0 \cdots i_n \cdots$, and any n

$$|\underline{\mathbf{i}}_2 - \underline{i_0 \cdots i_{n-1}}_2| \leq 2^{-n}$$

so, if n is large enough so that

$$|\underline{\mathbf{i}}_2 - \underline{\mathbf{i}'}_2| > 3 \times 2^{-n},$$

then

$$|\underline{i_0 \cdots i_{n-1}}_2 - \underline{i'_0 \cdots i'_{n-1}}_2| > 2^{-n},$$

which by Lemma 3.17 implies that $\overline{\mathbf{A}_{i_0 \cdots i_{n-1}}}$ and $\overline{\mathbf{A}_{i'_0 \cdots i'_{n-1}}}$ are disjoint, as asserted. ☐

Putting together what we know about the map $\mathbf{i} \mapsto \hat{z}(\mathbf{i})$, we finally obtain:

Proposition 3.19. *The angles* $\underline{\mathbf{i}}_2 = \underline{\mathbf{i}'}_2$ *if and only if* $\hat{z}(\mathbf{i}) = \hat{z}(\mathbf{i}')$. *Hence there is a function* $\theta \mapsto z(\theta)$ *from the circle* $\mathbb{R}/\mathbb{Z}$ *to* $\mathbb{C}$ *so that*

$$\hat{z}(\mathbf{i}) = \underline{\mathbf{i}}_2.$$

The function $z(\,.\,)$ *is continuous, bijective, and maps the circle onto*

$$J \equiv \cap_n (f_0^{-n} \overline{\mathbf{A}}).$$

We recall a well-known topological fact:

Theorem 3.20. *A continuous bijection from a compact to a Hausdorff topological space is, in fact, a homeomorphism.*

As a corollary, we have:

Corollary 3.21. *The set* J *is a Jordan curve.*

Note that

$$J \supset \cup_{n \geq 0} f^{-n}(0),$$

and hence $J \supset J(f_0)$. By construction, we have:

Proposition 3.22. *The map* $\underline{\mathbf{i}}_2 \mapsto \hat{z}(\mathbf{i})$ *is a conjugacy between* $\theta \mapsto 2\theta$ *and* $f_0(z)$*:*

$$f_0(\hat{z}(\mathbf{i})) = \hat{z}(2 \cdot \underline{\mathbf{i}}_2).$$

Hence, periodic orbits of f_0 are dense in J. Thus

$$J = J(f_0),$$

and the proof of Theorem 3.10 is completed.

3.4. **Covering properties of H_{f_0}.** We show:

Theorem 3.23. *The singular values of* H_{f_0} *are*

$$\mathrm{Sing}(H_{f_0}) = \{0, \infty, v\},$$

where $v = \mathrm{ixp} \circ \phi_A(-1/4)$*. The points* 0 *and* ∞ *are asymptotic values of* H_{f_0}*: there exists a path*

$$\gamma[0,1) \mapsto W^+ \text{ with } \gamma(t) \underset{t \to 1}{\longrightarrow} \partial W^+ \text{ and } f_0(\gamma(t)) \underset{t \to 1}{\longrightarrow} \partial 0,$$

and similarly for ∞*.*

Proof. Verifying that

$$H_{f_0}(\mathrm{Crit}(H_{f_0})) = \{v\}$$

is straightforward from the covering properties of the Fatou coordinate ϕ_A, and is left to the reader.

To fix the ideas, we show that the asymptotic values of the restriction $H_{f_0}|_{\hat{W}^+}$ are $\{0, \infty\}$; the argument for $H_{f_0}|_{\hat{W}^-}$ is identical. To simplify geometric considerations, let us invoke Proposition 3.25, and present the proof for $H \equiv H_h$ on a neighborhood of 0; normalized so that $H'(0) = 1$ and the domain of analyticity of H consists of the unit disk $\mathbb{D}$ and $\hat{\mathbb{C}} \setminus \overline{\mathbb{D}}$.

We demonstrate that

$$\operatorname{Asym}(H) = \{0, \infty\}.$$

Consider an attracting petal P for h which is bounded by a curve γ, which becomes a vertical line in Fatou coordinate ϕ^h_A, such that the critical value

$$-1/4 \in C \equiv P \setminus \overline{h(P)}.$$

It is easy to see that the second preimage $h^{-2}(\gamma)$ consists of a curve $\gamma_1 \subset \bar{\mathbb{H}}$ connecting the points 0 and 1, and γ_2 which does the same in the lower half-plane.

It is evident that $\{0, \infty\} \subset \operatorname{Asym}(H)$. Indeed, for small enough $\epsilon > 0$ the projection of $\gamma_1 \cap D_\epsilon(1)$ by $\operatorname{ixp} \circ \phi^h_R$ is a simple curve in $\mathbb{D}$ whose image under H lands at 0; similarly γ_2 yields a curve whose image under H lands at ∞.

Now let

$$\tau : [0, 1) \to \mathbb{D} = \operatorname{Dom}(H)$$

be a curve such that $\tau(t)$ converges to the unit circle $\mathbb{T}$ as $t \to 1-$. Let

$$\tilde{\tau} : [0, 1) \subset \hat{\mathbb{C}} \setminus \mathbb{R}_{\geq 0}$$

be a component of its lift under $\operatorname{ixp} \circ \phi^h_R$, parametrized so that

$$\tilde{\tau}(t) \to \mathbb{R}_{\geq 0} \text{ as } t \to 1- . \tag{3.5}$$

Let C' be a connected component of $h^n(C)$ for $n \in \mathbb{Z}$. We will say that the curve $\tilde{\tau}$ *crosses* C' if there exist $t_1 \neq t_2 \in (0, 1)$ and $n \in \mathbb{Z}$ such that

$$\tilde{\tau} : (t_1, t_2) \to C',\ h^n(\tilde{\tau}(t_1)) \in \gamma,\ \text{and } h^n(\tilde{\tau}(t_2)) \in h(\gamma).$$

Note that every time $\tilde{\tau}$ crosses some C', the image $H(\tau)$ winds once around the cylinder $\mathbb{C}/\mathbb{Z} \simeq \hat{\mathbb{C}} \setminus \{0, \infty\}$. Hence, we have the following two possibilities:

1. $\tilde{\tau}$ crosses finitely many connected components of $\cup_{n\in\mathbb{Z}} h^n(C)$;
2. $\lim_{t\to 1-} H(\tau(t)) \in \{0, \infty\}$.

In case (2), we are done.

Standard considerations of dynamics on $J(h)$ imply that for every $x \in \mathbb{R}_{\geq 0}$ which is not a preimage of 0, there exists an infinite sequence of cross-cuts $l_k \equiv h^{-n_k}(\gamma_i)$ for some choice of the inverse branch and $i = 1, 2$ such that denoting N_k the cross-cut neighborhood of l_k in $\hat{\mathbb{C}} \setminus \mathbb{R}_{\geq 0}$, we have

$$\overline{N_k} \cap \mathbb{R} \ni \{x\}.$$

In view of (1) and (2) this implies that x cannot be accumulated by $\tilde{\tau}$. Since points x as above form a dense set on $\mathbb{R}_{\geq 0}$, we see that there exists a limit

$$s = \lim_{t\to 1-} \tilde{\tau}(t) \text{ and } s \in \cup h^{-n}(0).$$

In view of (1), this implies that

$$\lim_{t\to 1-} H(\tau(t)) \in \{0, \infty\}.$$

□

3.5. **Parabolic renormalization of f_0.** We now have:

Proposition 3.24. *The maps f_0 and h are both renormalizable.*

Proof. The arguments are identical, so we only argue for f_0. Let us replace H_{f_0} by cH_{f_0} if needed so that $H'_{f_0}(0) = 1$. We have to show that the coefficient a in the Taylor expansion

$$H_{f_0}(z) = z + az^2 + \cdots$$

is not equal to 0, or, in other words, that $z = 0$ is a simple parabolic point of H_{f_0}. Let P_0 be an attracting petal of $z = 0$. For $n \geq 1$ denote P_{-n} the component of the preimage $(H_{f_0})^{-1}(P_{-(n-1)})$ which contains $P_{-(n-1)}$. We claim that there exists n such that P_{-n} contains the unique critical value v of H_{f_0}. Assume the contrary. Then for every n the domain P_{-n} is a topological disk. Denote

$$B_0 = \cup P_{-n} \subset \hat{W}^+.$$

The Fatou coordinate ϕ_A of H_{f_0} extends to all of B_0 via the functional equation

$$\phi_A \circ H_{f_0}(z) = \phi_A(z) + 1$$

as an unbranched analytic map. It is trivial to see that its image is the whole complex plane $\mathbb{C}$, and hence U must be a parabolic domain. This is impossible, however, as U is a subset of a Jordan domain $\hat{W}^+ \subset \mathbb{C}$.

Now assume that $a = 0$. Then the parabolic point $z = 0$ has at least two distinct attracting directions $v_1, v_2 \in S^1$. Consider corresponding attracting petals $P_A^1 \cap P_A^2 = \emptyset$. As we have shown above, the critical value v must be contained in the intersection of their preimages. This leads to a contradiction, as in this case

$$\lim H_{f_0}^n(v)/|H_{f_0}^n(v)| = v_1 = v_2.$$

□

Let us make a note:



FIGURE 1. The preimages of the real axis for the map h.

Proposition 3.25. *Denote $H \equiv H_h$ normalized so that*

$$H : \mathbb{D} \to \hat{\mathbb{C}}.$$

There exist $v \neq 0$ and $\theta \in \mathbb{R}$ such that

$$F \equiv \mathcal{P}(f_0) = v \cdot H \circ \psi(e^{2\pi i\theta} z), \text{ where } \psi : \mathbb{D} \to \hat{W}^+$$

is the conformal Riemann mapping, which satisfies $\psi(0) = 0$, $\psi'(0) > 0$.

We now turn to the dynamics of the parabolic renormalization

$$F \equiv \mathcal{P}(f_0) : \hat{W}^+ \to \hat{\mathbb{C}}.$$

Definition 3.3. The points $z \in \hat{W}^+$ whose orbits do not escape $\hat{W}^+$ form the *filled Julia set* $K(F)$. Its boundary $J(F)$ is the *Julia set*.

Lavaurs [Lav] has shown that

Theorem 3.26 ([Lav]). *The interior of the filled Julia set $K(F)$ coincides with the basin of the parabolic point 0. Repelling periodic orbits of F are contained in $J(F)$ and are dense in $J(F)$.*

Let $P^0 \equiv P_A$ be an attracting petal of F. For $n \in \mathbb{N}$ inductively define P^{-n} to be the connected component of the inverse image $F^{-1}(P^{-(n-1)})$ which contains $P^{-(n-1)}$. The same considerations as in the proof of Proposition 3.24 imply:

Proposition 3.27. *There exists $n \in \mathbb{N}$ such that P^{-n} is a topological disk which contains v in its closure.*

3.6. **A note on the general theory for analytic maps of finite type.** As the previous example clearly demonstrates, when f has global covering properties, we can expect the Écale-Voronin map $\mathcal{E}_f$ to also possess a well-understood global covering structure. The appropriate setting for a global structure theory for Écalle-Voronin maps is that of analytic maps *of finite type*, developed in A. Epstein's thesis [Ep]. While we will not need these results in our investigation, we will briefly mention some of them below for the sake of completeness of the exposition.

Definition 3.4 ([Ep]). Let $f : W \to X$ be an analytic map between two Riemann surfaces. Assume further that X is compact, and W lies in some compact surface Y. Suppose that f is nowhere constant, and that every isolated singularity of f is essential. We say that f is a map of finite type if $\mathrm{Sing}(f)$ is a finite set.

Rational endomorphisms $f : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ are obviously maps of finite type. Epstein demonstrated:

Theorem 3.28. [Ep] *If an analytic map f with a parabolic cycle is of finite type, then the corresponding Écalle-Voronin maps $\mathcal{E}_f : W_f \to \hat{\mathbb{C}}$ inherit the same property.*

Epstein further showed that the familiar properties, such as density of repelling periodic points and topological minimality, hold for Julia sets of analytic dynamical systems of finite type. Most importantly, he proved that the Fatou-Sullivan structure theory holds for the Fatou sets of such dynamical systems:

Theorem 3.29. [Ep] *Every connected component of the Fatou set of a map of finite type is pre-periodic. All periodic Fatou components are either basins (attracting or parabolic), or rotation domains (Siegel disks or Herman rings).*

4. A CLASS OF ANALYTIC MAPPINGS INVARIANT UNDER $\mathcal{P}$

4.1. **Definition of P.**

We will now specialize to a much narrower class of analytic maps with parabolic orbits. As before, we write

$$f_0(z) = z + z^2,$$

and we set

$$F \equiv \mathcal{P}(f_0) \text{ and } W_F = \mathrm{Dom}(\mathcal{P}(f_0)).$$

Definition 4.1. We will denote $\tilde{\mathbf{P}}$ the class of analytic germs $f(z)$ at the origin which have maximal analytic extensions

$$f : \mathrm{Dom}(f) \to \hat{\mathbb{C}}$$

such that

(I) $\mathrm{Dom}(f)$ is a Jordan domain;

(II) Denoting φ_f the Riemann mapping

$$\varphi_f : \mathbb{D} \to \mathrm{Dom}(f), \text{ with } \varphi_f(0) = 0 \text{ and } \varphi_f'(0) > 0,$$

we have

$$f \circ \varphi_f(z) = v \cdot F \circ \varphi_F(e^{2\pi i\theta} z), \text{ for } v \neq 0,\ \theta \in \mathbb{R}.$$

Note that denoting c and c^F the unique critical values of f and $F = \mathcal{P}(f_0)$ respectively, we get the rescaling factor as $v = c/c^F$.

If we again let $H \equiv H_h$ for $h(z)$ given by (3.2) normalized so that its maximal analytic extension about the origin has domain $\mathbb{D}$, then, by Proposition 3.25, the property (II) in the above definition is equivalent to

(II') Denoting φ_f the Riemann mapping

$$\varphi_f : \mathbb{D} \to \mathrm{Dom}(f), \text{ with } \varphi_f(0) = 0 \text{ and } \varphi_f'(0) > 0,$$

we have

$$f \circ \varphi_f(z) = v \cdot H(e^{2\pi i\theta} z), \text{ for } v \neq 0,\ \theta \in \mathbb{R}.$$

We note:

Proposition 4.1. *For all $f \in \tilde{\mathbf{P}}$ we have $f''(0) \neq 0$.*

Proof. Suppose $f''(0) = 0$. By multiplying by a non-zero constant, if necessary, we may reduce the proof to the case when $f'(0) = 1$, so that $f(z) = z + az^n + \cdots$ for $n > 2$. A Leau-Fatou flower of f has $n - 1 \geq 2$ attracting petals. Since f has a single critical value c^f, there exists an ample petal P_A such that

$$P_A \cap \cup_{k\in\mathbb{N}} f^k(c^f) = \emptyset.$$

Let ϕ_A be an attracting Fatou coordinate defined on P_A. Set $P_0 \equiv P_A$ and inductively define P_{-n} as the component of the preimage $f^{-1}(P_{-(n-1)})$ which contains $P_{-(n-1)}$ for $n \in \mathbb{N}$. The function ϕ_A analytically extends via the functional equation

$$\phi_A \circ f(z) = \phi_A(z) + 1$$

to the union

$$U \equiv \cup P_{-n} \subset \mathrm{Dom}(f).$$

A simple induction shows that ϕ_A is univalent on P_{-n}, and hence on all of U. On the other hand, $\phi_A(U) = \mathbb{C}$, which is impossible since U is a hyperbolic domain. □

Definition 4.2. We define $\mathbf{P}$ as the set of maps $f \in \tilde{\mathbf{P}}$ of the form

$$f(z) = z + z^2 + \cdots$$

at the origin.

Since $F = \mathcal{P}(f_0) \in \mathbf{P}$, we have

Proposition 4.2. *The class* $\mathbf{P}$ *is non-empty.*

We now set out to prove the following theorem:

Theorem 4.3. *Every* $f \in \mathbf{P}$ *is renormalizable, and the parabolic renormalization* $\mathcal{P}(f) \in \mathbf{P}$.

Let $f \in \mathbf{P}$ and let P_0 be an attracting petal of the parabolic point $z = 0$. For $n \geq 1$ let P_{-n} be the component of $f^{-1}(P_{-(n-1)})$ which contains $P_{-(n-1)}$. We let

$$B_0^f \equiv \cup P_{-n},$$

and call it the *immediate basin* of 0. We note:

Proposition 4.4. *For* $f \in \mathbf{P}$ *the unique critical value* $c^f \in B_0^f$. *Moreover, there exists an attracting petal* $P_A \subset B_0^f$ *which is a topological disk containing* c^f *in its interior.*

Proof. Assume that $v \notin B_0^f$. The attracting Fatou coordinate ϕ_A^f extends from P_0 to the whole of B_0^f via the functional equation

$$\phi_A^f \circ f(z) = \phi_A^f(z) + 1.$$

A simple induction shows that P_{-n} is an increasing sequence of topological disks, on each of which ϕ_A is univalent. Hence, ϕ_A restricted to B_0^f is a conformal homeomorphism onto the image. Yet it is clear that the image of ϕ_A restricted to B_0^f is the whole of $\mathbb{C}$, which contradicts the fact that B_0^f is a hyperbolic domain. The second part of the statement is elementary, and is left to the reader. □

Pushing the argument a little further, we have:

Proposition 4.5. *For $f \in \mathbf{P}$ the immediate basin B_0^f is simply connected and contains exactly one critical point of f. The restriction $f : B_0^f \to B_0^f$ is a degree-2 branched covering.*

Proof. By Proposition 4.4, there exists n such that P_{-n} contains the critical value c^f. Inductively applying Lemma 3.7, we see that for $k \in \mathbb{N}$ the domain $P_{-(n+k)}$ is a topological disk, and $f : P_{-(n+k)} \to P_{-(n+k-1)}$ is a branched covering of degree 2 with a single, simple critical point. $\square$

The proof of Theorem 4.3 will rely on the following key result, which is a direct analogue of Theorem 3.10.

Theorem 4.6. *Let $f \in \mathbf{P}$, and denote B_f^0 the immediate basin of the parabolic point 0 of f. Then B_f^0 is a Jordan domain. Denote $\hat{f}$ the continuous extension of f to ∂B_0^f. There exists a homeomorphism $\rho : \partial B_0^f \to \mathbb{T}$ such that*

$$\rho(\hat{f}(z)) = 2\rho(z) \bmod 1.$$

The proof of Theorem 4.6 is quite involved, as it will require a detailed understanding of the covering properties of a map in $\mathbf{P}$. Let us show how Theorem 4.6 implies Theorem 4.3:

Proof. Consider a Riemann map

$$v : \mathbb{D} \to B_0^f$$

which maps 0 to the sole critical point of f inside B_0^f. By Theorem 4.6 and Carathéodory Theorem, the Riemann map has a continuous extension to a homeomorphism $\tilde{v} : \overline{\mathbb{D}} \to \overline{B_0^f}$.

Let us further normalize the Riemann map $\mathbb{D} \to B_0^f$ by requiring that $\tilde{v}(1) = 0$. This specifies the mapping uniquely, and we denote it ψ_f.

Set

$$B(z) = (\psi_f)^{-1} \circ f \circ \psi_f : \mathbb{D} \to \mathbb{D}.$$

The map B is a degree two Blaschke product, with a single simple ramification point at the origin. It fixes the boundary point 1. By the local dynamics of f, the point 1 attracts some orbits of B in $\bar{\mathbb{D}}$, and hence $|B'(1)| \leq 1$.

Assume that $|B'(1)| < 1$. Then there exists an arbitrarily small crosscut γ forming a crosscut neighborhood N of the boundary point $p = 0$ in B_0^f, whose forward orbit is entirely contained in N. This clearly contradicts the local dynamics of the parabolic point $p = 0$, since the boundary points $\partial\gamma \subset \partial B_0^f$ lie in a repelling petal of f.

Hence $|B'(1)| = 1$, which evidently means $B'(1) = 1$. Applying Schwarz reflection about the circle we see that the parabolic point 1 has at least two attracting directions, and hence $B''(1) = 0$.

An inspection shows that the unique Blaschke product of degree two with the properties $B'(0) = 0$, $B(1) = 1$, $B'(1) = 1$, $B''(1) = 0$ is given by the formula

$$B(z) = \frac{3z^2 + 1}{3 + z^2}.$$

Let us set

$$\chi = \psi_{f_0} \circ (\psi_f)^{-1} : B_0^f \mapsto B_0^{f_0}.$$

By the discussion above, this mapping is a conjugacy:

$$\chi \circ f|_{B_0} = f_0 \circ \chi|_{B_0}.$$

Let us fix

$$D = \{|w| < |c^f|\},$$

and let g be the branch of f^{-1} fixing 0 which is defined in D. For the moment, the map $\mathcal{P}_f$ is only defined locally in a neighborhood of 0, which is *a priori* dependent on the choice of a repelling petal P_R^f. Let us fix a petal $P_R^f \subset D$.

The map $F = \mathcal{P}(f_0)$ has a maximal extension to a Jordan domain W_F. Let us use Theorem 3.10 to select a simple arc γ connecting 0 with $a \in J_{f_0}$ inside $B_0^{f_0} \cap P_R^{f_0}$ in some repelling petal of f_0, such that:

- $\gamma \cap J_{f_0} = \{a\}$;
- $\gamma \cap f_0(\gamma) = \{0\}$;
- denoting $V_0 \subset P_R^{f_0}$ the Jordan domain bounded by γ, $f_0(\gamma)$, and the Jordan subarc of J_{f_0} between a and $f_0(a)$, we have a one-to-one correspondence between points in W_{f_0} and points in
$$\hat{V}_0 \equiv (V_0 \cup \gamma) \setminus \{a\};$$
- $\hat{V} \equiv \chi(\hat{V}_0) \subset P_R^f$.

Denote $\mathcal{W} \subset \mathcal{C}_R^f$ the quotient of $\hat{V}$ by the action of f. Let $W \subset \mathbb{C}$ be the image of $\mathcal{W}$ by $\operatorname{ixp} \circ \phi_R^f$ with the puncture at 0 filled. By construction, W is a Jordan domain, which contains a neighborhood of 0. We claim that:

(1) the germ $\mathcal{P}(f)$ analytically continues to all of W, and, moreover,
(2) $\mathcal{P}(f)$ cannot be analytically continued to a neighborhood of any point $z \in \partial W$.

Since $V \subset P_R^f \cap B_0^f$, the first statement is immediate by the definition of $\mathcal{P}(f)$. Furthermore, by (3.1), the set of critical points

$$\operatorname{Crit}(\phi_A^f) = \chi(\operatorname{Crit}(\phi_A^{f_0}))$$

contains ∂B_0^f in its closure. Since ∂W is a projection of a Jordan subarc of ∂B_0^f, this implies (2). We have thus shown the existence of a Jordan domain

$$\mathrm{Dom}(f) = W.$$

Let

$$\varphi_f : \mathrm{Dom}(f) \to \mathbb{D}$$

be the unique Riemann mapping normalized as in Definition (4.1). The uniqueness of the Riemann mapping implies that, up to a precomposition with a rotation, it is given by the composition

$$\mathrm{ixp} \circ \phi_R^f \circ \chi^{-1} \circ (\phi_R^{f_0})^{-1} \circ \mathrm{ixp}^{-1} \circ \varphi_F,$$

with suitably chosen inverse branches.

The map χ induces a conformal isomorphism between attracting Fatou cylinders $\mathcal{C}_A^f \to \mathcal{C}_A^{f_0}$. After uniformizing the cylinders by $\mathbb{C}^*$, it takes the form

$$\tau \equiv \mathrm{ixp} \circ \phi_A^{f_0} \circ \chi \circ (\phi_A^f)^{-1} \circ \mathrm{ixp}^{-1} : \mathbb{C}^* \to \mathbb{C}^*$$

(with suitably chosen inverse branches). By Proposition 2.10, τ is a multiplication by a non-zero constant.

A diagram chase now implies that

$$\mathcal{P}_f \circ \varphi_f(z) = v \cdot F \circ \varphi_F(e^{2\pi i\theta} z)$$

for some $v \neq 0$ and $\theta \in \mathbb{R}$. Since the unique critical value of the mapping on the left coincides with that of the mapping on the right, $v = c^{\mathcal{P}(f)}/c^F$.

□

Let $f \in \mathbf{P}$. By definition of parabolic renormalization, for any repelling petal P_R^f the projection of the intersection $P_R^f \cap B_0^f$ by $\mathrm{ixp} \circ \phi_R^f$ lies in the domain $\mathrm{Dom}(\mathcal{P}(f))$. We remark, that it covers all of it:

Remark 4.1. For any choice of P_R^f, the projection

$$\mathrm{ixp} \circ \phi_R^f(P_R^f \cap B_0^f) \cup \{0, \infty\}$$

is a union of two disjoint Jordan domains $W^+ \ni 0$ and $W^- \ni \infty$ with

$$W^+ = \mathrm{Dom}(\mathcal{P}(f)).$$

We begin the proof with a lemma:

Lemma 4.7. *Let D be a Jordan domain containing* 0 *such that*

- $c^f \notin D$;
- *there exists a Jordan arc $\tau \ni c^f$ in $\hat{\mathbb{C}} \setminus D$ running to ∞ such that $\tau \cap B_0^f$ is connected.*

Analytically extend g *to* D*. If* $w \in B_0^f \cap D$*, then* $g(w) \in B_0^f$*.*

Proof. Standard considerations imply that g extends to a branch $\hat{g}$ of f^{-1} defined and analytic on $\mathbb{C} \setminus \tau$. Note that $B_0^f \setminus \tau$ is also connected. For a point z_0 on the negative real axis and near to 0, the asymptotic development for the attracting Fatou coordinate (Proposition 2.5) implies that z_0 and $g(z_0) \approx z_0$ both lie in B_0^f.

Let $z \in B_0^f \setminus \tau$; then there is a Jordan arc γ in $B_0^f \setminus \tau$ from z_0 to z. By the covering properties of f, there is a unique lift $\tilde{\gamma}$ of γ starting at $g(z_0)$. Furthermore, this lift is contained in B_0^f, so its end point is in B_0^f. On the other hand, for $\hat{g}$ as above,

$$s \mapsto \hat{g}(\gamma(s))$$

is another lift with the same starting point. By uniqueness of lifts, it coincides with $\tilde{\gamma}$. In particular, the end point of $\tilde{\gamma}$ is $\hat{g}(z)$. Since we already know that $\tilde{\gamma}$ lies in B_0^f, it follows that $\hat{g}(z) \in B_0^f$. In particular, if $z \in D$, then $\hat{g}(z) = g(z)$, so $g(z) \in B_0^f$, as asserted. □

Proof of Remark 4.1. Extend the local inverse g to all of P_R^f. If necessary, replace P_R^f with $g^n(P_R^f)$ for a sufficienly large $n \in \mathbb{N}$ to guarantee that there exists a domain D as described in Lemma 4.7 such that $P_R^f \subset D$.

Denote

$$W = \operatorname{ixp} \circ \phi_R^f(P_R^f \cap B_0^f) \text{ and } V = \operatorname{ixp} \circ \phi_R^f(P_R^f \setminus B_0^f).$$

Lemma 4.7 implies that W and V are disjoint. The common boundary $J = \partial W = \partial V$ is the projection

$$J = \operatorname{ixp} \circ \phi_R^f(P_R^f \cap \partial B_0^f).$$

By Theorem 4.6, J is a union of two disjoint Jordan curves. Hence, W is a union of two Jordan domains $W^+ \ni 0$ and $W^- \ni \infty$, bounded by the components of J. On the other hand, $\operatorname{Dom}(\mathcal{P}(f))$ is also a Jordan domain, which does not intersect J. Hence, it is contained in the component of W which surrounds 0, and is, in fact, equal to it by the maximality of the analytic continuation of $\mathcal{P}(f)$. □

4.2. **The structure of the immediate parabolic basin of a map in H.** In this section we will prove several results about lifts of parametrized paths of the form

$$s : [0, 1] \to \hat{\mathbb{C}}.$$

We will always assume $s(t)$ to be continuous on $(0,1)$. We will say that s *lands* at a point $a \in \hat{\mathbb{C}}$ if

$$\lim_{t\to b} s(t) = a \text{ for } b \in \{0,1\}.$$

We will generally use the same letter s to denote the function $s(t)$ and the curve $s([0,1])$ which is its range. We will call the image of the open interval $(0,1)$ under $s(t)$ an *open* path, and will denote it by $\overset{\circ}{s}$. Several times we will encounter the situation when there is a domain $W \subset \hat{\mathbb{C}}$ and a curve

$$s : (0,1) \mapsto W,$$

which lands at a point $w \in \partial W$ (to fix the ideas, assume that $s(0) = w$). If w has more than one prime end in W, then there is a unique prime end $\hat{w}$ such that for every prime end neighborhood $N(\hat{w})$ we have $s \cap N(\hat{w}) \neq \emptyset$. In this case, we will write

$$s(t) \underset{t\to 0+}{\longrightarrow} \hat{w}.$$

Let $f \in \mathbf{P}$. Let $s \mapsto \gamma(s)$ be a continuous path such that:

- $\gamma(0) = \gamma(1) = 0$;
- $\gamma(s) \neq c^f$ for all $s \in (0,1)$;
- the winding number $W(\gamma, c^f) = 1$;
- there is an $\epsilon \in (0, \pi/2)$ such that $\gamma \cap D_\epsilon(0)$ lies in the sector $\{\mathrm{Arg}(z) \in (-\pi - \epsilon, -\pi + \epsilon)\}$.

Standard path-lifting considerations imply that there exists a unique continuous mapping $s \mapsto \tilde{\gamma}(s)$, defined for $0 \leq s < 1$ such that

$$\tilde{\gamma}(0) = 0, \text{ and } f(\tilde{\gamma}(s)) = \gamma(s) \text{ for } 0 \leq s < 1.$$

We claim:

Proposition 4.8. *We have the following.*

(1) *For any loop γ as above, there exists a limit*

$$t \in \partial\,\mathrm{Dom}(f) = \lim_{s\to 1} \tilde{\gamma}(s).$$

(2) *This point (which we will denote t^f) is the same for all loops γ satisfying the above properties.*

(3) *For $f = F \equiv \mathcal{P}(f_0)$, the point t^F is the projection of the inverse orbit*

$$z_{-n} = (g_0)^n(-1)$$

by $\mathrm{ixp} \circ \phi_R$.



FIGURE 2. An illustration of the proof of Proposition 4.8

(4) *Similarly, for* $f = H \equiv \mathcal{P}(h)$*, the point* t^H *is the projection of the inverse orbit*

$$z_{-n} = h^{-n}(1),$$

where the inverse branch is selected to preserve the interval $(0, 1)$.

Proof of Proposition 4.8. We will prove (1) - (3) for $f = F \equiv \mathcal{P}(f_0)$. By the definition of $\mathbf{P}$ this will imply (1) and (2) in the general case. The proof of (4) follows along the same lines as the proof of (3) and will be left to the reader.

Let us begin by selecting a simple curve $\sigma \subset \hat{\mathbb{C}}$ which connects 0 and ∞, does not intersect with γ except at the endpoint 0, and such that an analytic branch of the logarithm defined on a neighborhood of σ has bounded imaginary part.

Recall that the attracting Fatou coordinate $\phi_A^{f_0}$ holomorphically extends to the whole immediate basin $B_0^{f_0}$ via the equation

$$\phi_A^{f_0} \circ f_0(z) = \phi_A^{f_0}(z) + 1.$$

This extension is a branched covering $B_0^{f_0} \to \mathbb{C}$ with simple ramification points at preimages of the critical point $-1/2$.

It is elementary to see what the lift of σ to the dynamical plane of f_0 looks like. We summarize its properties below, and invite the reader to verify them. Let us denote

$$\overset{\circ}{\sigma} \equiv \sigma \setminus \{0, \infty\}.$$

The preimage of $\overset{\circ}{\sigma}$ under $\text{ixp} \circ \phi_A^{f_0}$ is a countable collection of disjoint simple curves $\cup \overset{\circ}{s}_j$. We will denote the closure of $\overset{\circ}{s}_j$ by s_j. It is obtained by adjoining two endpoints to $\overset{\circ}{s}_j$: two elements of the grand orbit $\cup_{n\geq 0}(f_0)^{-n}(0)$, which are not necessarily distinct.

The curves s_j form a grand orbit under f_0 as well. The connected components

$$S_i \text{ of } B_0^{f_0} \setminus \cup s_j$$

are mapped by the attracting Fatou coordinate $\phi_A^{f_0}$ onto curvilinear strips $\tilde{S}_i \subset \mathbb{C}$ of infinite height and of unit width. The strips $\tilde{S}_i$ are bounded by unit translates of the same curve $\tilde{\sigma} \subset \mathbb{C}$. Let us enumerate these strips in such a way that

$$T : \tilde{S}_i \to \tilde{S}_{i+1}, \text{ where } T(z) = z + 1.$$

For a fixed $\tilde{S}_i$ we have

$$\sup_{x,y \in \tilde{S}_i} (\text{Re}(x) - \text{Re}(y)) < \infty.$$

This is where we have used the assumption on the branches of log in $\hat{\mathbb{C}} \setminus \sigma$. Thus for every $\ell \in \mathbb{Z}$ the union $\cup_{i \geq \ell} \tilde{S}_i$ contains a right half-plane.

Now let us take *any* component $S_i \subset B_0^{f_0}$. The above observation implies that, for any $\ell \in \mathbb{N}$,

$$\phi_A^{f_0}(\cup_{k\geq \ell} f^k(S_i)) \supset \{\text{Re}\, z > A\}$$

for some $A \in \mathbb{R}$. Note that if ℓ is large enough, then $\cup_{k\geq\ell} f^k(S_i)$ does not contain any preimages of $-1/2$, and hence the restriction of $\phi_A^{f_0}$ to it is unbranched. Denote the boundary components of $S_{i+\ell}$ by s and $f_0(s)$. Then s bounds an attracting petail P, and

$$P = \cup_{k\geq\ell} f^k(S_i).$$

We now complete the proof as follows. Fix

$$S \equiv S_{i+\ell} = P \setminus \overline{f_0(P)}.$$

The curve $\mathring{\gamma} = \gamma((0,1))$ has a univalent pull-back $\mathring{\nu} \subset S$. We continuously extend it to a parametrized loop

$$\nu : [0,1] \to S \cup \{0\}$$

which starts and ends at the parabolic point 0.

Denote $P_0 \equiv P$ and, for $j \in \mathbb{N}$, let P_{-j} be the connected component of $(f_0)^{-1}(P_{-(j-1)})$ which contains $P_{-(j-1)}$. By Proposition 4.4, there exists $n \in \mathbb{N}$ such that P_{-n} is an attracting petal such that the critical value $-1/4$ is contained in $P_{-n} \setminus \overline{P_{-(n-1)}}$. Denote $\nu_0 = \nu$ and for $1 \leq j \leq n$ let $\nu_{-j} \subset P_{-j} \setminus P_{-(j-1)}$ be the univalent pull-back of $\nu_{-(j-1)}$. The curve ν_{-n} is a parametrized loop

$$\nu_{-n} : [0,1] \to B_0^{f_0} \text{ with } \nu_{-n}(0) = \nu_{-n}(1) = 0.$$

By the choice of n, the winding number

$$W(\nu_{-n}, -1/4) = 1.$$

Consider the parameterized curve

$$\nu_{-(n+1)} \subset P_{-(n+1)} \setminus P_{-n}$$

such that $\nu_{-(n+1)}(0) = 0$ and

$$f_0(\nu_{-(n+1)}) = \nu_{-n}.$$

Evidently,

$$\nu_{-(n+1)}(1) = -1,$$

which is the *other* preimage of the parabolic point $z = 0$ under f_0. To complete the argument, let us now consider the connected component W of

$$B_0^{f_0} \setminus \overline{P_{-(n+1)}} \supset W$$

which contains the "upper" preimage of -1, the point

$$z_{-1} \equiv \frac{-1+\sqrt{3}i}{2} \in \mathbb{H},$$

in the boundary. Let $\nu_{-(n+2)} \subset W$ be the next preimage,

$$f_0(\nu_{-(n+2)}) = \nu_{-(n+1)}, \text{ with } \nu_{-(n+2)}(0) = 0 \text{ and } \nu_{-(n+2)}(1) = z_{-1}.$$

The inverse branch g_0 of the quadratic map f_0 univalently extends to a map

$$g_0 : W \to W.$$

For $k \geq 3$ denote $\nu_{-(n+k)}$ the preimage of $\nu_{-(n+k-1)}$ by this branch. By Denjoy-Wolff Theorem,

$$\nu_{-(n+k)} \to 0.$$

Furthermore, $\nu_{-(n+k)}$ is disjoint from $P_{-(n+1)}$. For any given ample repelling petal P_R there exists k such that $\nu_{-(n+k)} \subset P_R$. Consider the projection

$$\tilde{\gamma} \equiv \operatorname{ixp} \circ \phi_R^{f_0}(\nu_{-(n+k)}).$$

By construction, it satisfies the properties (1)-(3). □

To help understand the shape of the immediate basin of $F \in \mathbf{P}$, let us look at the drawing in Figure 4.2. It illustrates the case $F = \mathcal{P}(f)$ (the reader may think of $f = f_0$, to fix the ideas). The left figure shows a fragment of the boundary of the immediate basin B_0^f near the parabolic fixed point 0. If we look on the right, we see a schematic picture of the Jordan domain $\operatorname{Dom}(\mathcal{P}(f))$. The point $t^{\mathcal{P}(f)} \in \partial \operatorname{Dom}(\mathcal{P}(f))$ is the "tip of the tail" of the immediate basin $B_0^{\mathcal{P}(f)}$. On the left the reader can see how the tail is formed. The lift of the basin B_0^f fits inside a suitable repelling crescent C_R^f, reaching to the upper tip of the crescent (which under $\operatorname{ixp} \circ \phi_R^f$ becomes the parabolic point $z = 0$ on the right). The point $t^{\mathcal{P}(f)}$ lifts under $\operatorname{ixp} \circ \phi_R^f$ to an f^{-k}-preimage $w \in C_R^f$ of the parabolic point 0. The lift of the immediate basin $B_0^{\mathcal{P}(f)}$ to C_R^f reaches to w; its shape near w is a conformal image of the shape near 0 (a tail). The map $\operatorname{ixp} \circ \phi_R^f$ is conformal in a neighborhood of w, and hence the basin $B_0^{\mathcal{P}(f)}$ also has a tail ending at $t^{\mathcal{P}(f)}$.

Let us make a new definition. Let $f \in \mathbf{P}$. Consider an attracting petal P_A^f which contains the unique critical value $c^f \in B_0^f$. Let $\tau \subset B_0^f$ be a simple curve which connects c^f with the parabolic point 0 and has the property $f(\tau) \subset \tau$. To fix the ideas, we will take this curve to be the horizontal ray

$$\{\operatorname{Re}(z) \geq \operatorname{Re}(\phi_A^f(c^f)),\ \operatorname{Im}(z) = \operatorname{Im}(\phi_A^f(c^f))\}$$

in the Fatou coordinate. Using Proposition 4.8, and passing from a curve around the critical value to a slit connecting the critical value with a parabolic point we see:

Proposition 4.9. *There exists a unique simple curve Γ^{in} connecting t^f with 0 such that*

$$f(\Gamma^{in}) = \tau.$$

Let us use the fact that $\operatorname{Dom}(f)$ is a Jordan domain, and hence the tip of the tail $t^f \in \partial \operatorname{Dom}(f)$ is accessible from outside $\operatorname{Dom}(f)$, and let us continue Γ^{in} by attaching a simple curve Γ^{out} connecting t^f to ∞



FIGURE 3. The formation of a "tail" in the immediate parabolic basin of $\mathcal{P}(f)$.

without intersecting $\overline{\mathrm{Dom}(f)}$. Let us further require that Γ^{out} coincides with a negative real ray in a neighborhood of ∞.

Definition 4.3. We call $\Gamma \equiv \overline{\Gamma^{\mathrm{in}} \cup \Gamma^{\mathrm{out}}}$ the *primary cut* of f.

We prove the following:

Proposition 4.10. *Let $f \in \mathbf{P}$. There exists a domain $U \subset \mathbb{C}$ such that:*

(1) *U is a Jordan domain;*
(2) *$U \subset \mathrm{Dom}(f)$ and $\partial U \cup \partial\,\mathrm{Dom}(f) = \{t^f\}$;*
(3) *$\partial U \subset f^{-1}(\overline{\Gamma^{out}} \cup \partial\,\mathrm{Dom}(f))$ and $U \supset \Gamma^{in}$;*
(4) *$U \ni c^f$;*
(5) *there is a single critical point $p^f \in \Gamma^{in}$ of f inside U.*

Finally, there is a simple arc $\Gamma^{in}_{-1} \subset f^{-1}(\Gamma^{in}) \cap U$ with endpoints $u^+, u^- \in \partial U$ such that $\Gamma^{in}_{-1} \cap \Gamma^{in} = \{p^f\}$ for which the following holds:

(6) *consider the Jordan domain U' which is the connected component of $U \setminus \Gamma^{in}_{-1}$ whose boundary does not contain the point t^f. Then*
$$f : U' \setminus \Gamma^{in} \longrightarrow \mathrm{Dom}(f) \setminus \Gamma^{in}$$
is a univalent map.

In view of the definition of $\mathbf{P}$, it suffices to prove a slightly more general statement for $f = F \equiv \mathcal{P}(f_0)$:



$f_0(a_{-(n+k)})$
These points are identified on the repelling Fatou cylinder
f_0^k
$a_{-(n+k)}$
$\nu_{-(n+k)}$
$S_{-(n+k)}$
0
ν_{-n}
S_{-n}
A pull-back of Γ^{out}
A pull-back of the boundary of Dom(F)
B_0^F
Γ^{out}
Dom(F)
Γ^{in}

Figure 4. An illustration to the proof of Proposition 4.11. The pull-back of $\hat{\mathbb{C}} \setminus \Gamma$ for $F = \mathcal{P}(f_0)$.

Proposition 4.11. *Let* $F = \mathcal{P}(f_0)$. *Consider any simple arc* $\gamma = \gamma_1 \cup \gamma_2$ *and any Jordan curve* τ *with the following properties:*

- γ_1 *is a simple arc which connects* 0 *and* t^F*;*
- γ_2 *is a simple arc which connects* t^F*;*
- *the curve* γ_1 *approaches* 0 *within a sector* $\{\operatorname{Arg}(z) \in (\pi/2, 3\pi/2)\}$*;*
- *the curve* γ_2 *approaches* ∞ *within a sector* $\{\operatorname{Arg}(1/z) \in (\pi/2, 3\pi/2)\}$*;*
- *the curve* τ *separates* c^F *from* ∞ *and* $\tau \cap \gamma = \{t^F\}$.

There exists a domain U *of* $F^{-1}(V)$ *such that:*

1. U *is a Jordan domain;*
2. $U \subset \operatorname{Dom}(F)$ *and* $\partial U \cup \partial \operatorname{Dom}(F) = t^F$*;*
3. $\partial U \subset F^{-1}(\overline{\gamma_2} \cup \tau)$;
4. U *contains a single critical point* p^F*;*

(5) *there is a simple arc* $\delta \subset U$ *which is a preimage of* γ_1, *has endpoints on the boundary of* U, *and intersects* γ_1 *at the critical point* p^F *only. If we let* U' *be the component of* $U \setminus \delta$ *not containing* t^F *on the boundary, then the map*

$$F : U' \longrightarrow \mathrm{Dom}(F) \setminus F(\delta)$$

is univalent.

Proof. Let us again begin by selecting a simple curve $\sigma \subset \hat{\mathbb{C}}$ which connects 0 and ∞, and does not intersect with $\gamma \cup \tau$ except at the point 0, and such that an analytic branch of the logarithm defined on a neighborhood of σ has bounded imaginary part.

Let us parametrize γ:

$$\gamma(t) : [0,1] \to \hat{\mathbb{C}}$$

so that $\gamma(0) = 0$ and $\gamma(1) = \infty$.

With a slight abuse of notation we use the same notation for the preimages of σ and γ, etc. as in the proof of Proposition 4.8. Let S_{-n} be the fundamental crescent $P_{-n} \setminus \overline{f(P_{-n})}$ which contains the critical value $-1/4$. The corresponding component $\nu_{-n} \subset \overline{S_{-n}}$ of the lift of γ is a simple arc which passes through $-1/4$, and whose two ends land at 0.

There are two prime ends of the point 0 in the crescent S_{-n}. We denote the “upper” one by 0^+_{-n}, and the “lower” one by 0^-_{-n}. They correspond to the points 0 and ∞ respectively in the quotient $\mathcal{C}_A \simeq \hat{\mathbb{C}}$. We will write

$$\nu_{-n}(t) \underset{t\to 0+}{\longrightarrow} 0^+_{-n} \text{ and } \nu_{-n}(t) \underset{t\to 1-}{\longrightarrow} 0^-_{-n}.$$

Writing $S_{-(n+k)} = P_{-(n+k)} \setminus P_{-(n+k-1)}$, we see that

$$f_0 : S_{-(n+1)} \to S_{-n}$$

is a double covering, branched at

$$-1/2 \mapsto -1/4 \in \nu_{-n}.$$

We thus obtain two parametrized curves $\nu^1_{-(n+1)}$ and $\nu^2_{-(n+1)}$ as the lifts of ν_{-n} by f_0. The intersection

$$\overset{\circ}{\nu}{}^1_{-(n+1)} \cap \overset{\circ}{\nu}{}^2_{-(n+1)} = -1/2.$$

For $j \leq n+1$ we denote 0^+_{-j}, 0^-_{-j} the “upper” and the “lower” prime ends of 0 in S_{-j}, so that

$$\tilde{f}_0 : 0^\pm_{-j} \mapsto 0^\pm_{-(j-1)}.$$

The other preimage of 0 in $\partial S_{-(n+1)}$ is the point -1. We use $-1^+_{-(n+1)}$ and $-1^-_{-(n+1)}$ to denote its two prime ends in $S_{-(n+1)}$ labeled so that the Carathéodory extension $\tilde{f}_0$ maps

$$-1^+_{-(n+1)} \mapsto 0^+_{-n} \text{ and } -1^-_{-(n+1)} \mapsto 0^-_{-n}.$$

We have

$$\nu^1_{-(n+1)}(t) \underset{t\to 0+}{\longrightarrow} 0^+_{-(n+1)} \text{ and } \nu_{-(n+1)}(t) \underset{t\to 1-}{\longrightarrow} -1^-_{-(n+1)};$$

$$\nu^2_{-(n+1)}(t) \underset{t\to 0+}{\longrightarrow} -1^+_{-(n+1)} \text{ and } \nu_{-(n+1)}(t) \underset{t\to 1-}{\longrightarrow} 0^-_{-(n+1)}.$$

Denote $S_{-(n+k)} \subset W$ the pull-back of $S_{-(n+k-1)}$ by the inverse branch g_0; and

$$\nu^{1,2}_{-(n+k)} \subset S_{-(n+k)}$$

the corresponding preimage of ν_{-n}. A trivial induction shows that there are exactly three points

$$\{0, a_{-(n+k)}, f(a_{-(n+k)})\} \subset \partial S_{-(n+k)},$$

which map to 0 under f_0^k. Furthermore, $a_{-(n+k)}$ has two prime ends in $S_{-(n+k)}$; the points 0 and $f_0(a_{-(n+k)})$ each have a single prime end. We denote $a^{\pm}_{-(n+k)}$ the prime end mapped to $0^{\pm}_{-n}$ by the Carathéodory extension $\tilde{f}_0^k$.

We have

$$\nu^1_{-(n+k)}(t) \underset{t\to 0+}{\longrightarrow} 0_{-(n+k)} \text{ and } \nu_{-(n+1)}(t) \underset{t\to 1-}{\longrightarrow} f_0(a_{-(n-k)});$$

$$\nu^2_{-(n+k)}(t) \underset{t\to 0+}{\longrightarrow} a^+_{-(n+1)} \text{ and } \nu_{-(n+k)}(t) \underset{t\to 1-}{\longrightarrow} a^-_{-(n+1)}.$$

Each of the sets ν_{-j} is composed of two parts, $\nu_{-j,1}$ and $\nu_{-j,2}$ which are the preimages of γ_1 and γ_2 respectively. For $j \leq n$, they intersect at a single point b_{-j}; for $j > n$ there are two such points b^1_{-j}, b^2_{-j}.

Denote $\kappa_{-(n+k)} \subset S_{-(n+k)}$ the preimage of τ. Elementary considerations of monodromy, which we spare the reader, imply that $\kappa_{-(n+k)}$ is a union of two simple arcs: one connecting 0 with $b^1_{-(n+k)}$, the other with $b^2_{-(n+k)}$; and otherwise disjoint from $\nu_{-(n+k)}$.

Considerations of Denjoy-Wolff Theorem imply that for every ample repelling petal P_R there exists k such that $S_{-(n+k)} \subset P_R$. Let us project the picture in $S_{-(n+k)}$ back to $\hat{\mathbb{C}}$ using $\operatorname{ixp} \circ \phi_R^{f_0}$. The points $a_{-(n+k)}$ and $f_0(a_{-(n+k)})$ are identified in the projection. We obtain a Jordan domain U, bounded by the projections of $\kappa^1_{-(n+k)}$, $\kappa^2_{-(n+k)}$, $\nu_{-(n+k),2}$. It satisfies the properties (1)-(5) by the construction.

□

Corollary 4.12. *Let $f \in \mathbf{P}$, and let U be the domain constructed in Proposition 4.10. We have*

- $B_0^f \subset U$;
- $\partial B_0^f \cap \partial \operatorname{Dom}(f) = \{t^f\}$.

Proof. Since B_0^f does not intersect with $\overline{\Gamma^{\text{out}}} \cup \partial \operatorname{Dom}(f)$ and $\partial U \subset f^{-1}(\overline{\Gamma^{\text{out}}} \cup \partial \operatorname{Dom}(f))$, we have $B_0^f \subset U$. Hence,

$$\partial B_0^f \cap \partial \operatorname{Dom}(f) \subset \{t^f\}.$$

On the other hand, the cut $\Gamma^{\text{in}} \subset B_0^f$, hence

$$t^f \in \overline{\Gamma^{\text{in}}} \in \partial B_0^f.$$

□

4.3. **The immediate basin of a map in P is a Jordan domain.** We are now in a position to begin the proof of Theorem 4.6. Let us fix $F \in \mathbf{P}$ and let U be the domain constructed in Proposition 4.10. The preimage $\overline{F^{-1}(\Gamma^{\text{in}})} \cap B_0^F$ consists of the curve Γ^{in} and a simple curve Γ^{in}_{-1}; the two curves cross at the critical point $p^F \in B_0^F$. To fix the ideas, let us parametrize

$$\Gamma^{\text{in}} : [0,1] \to \overline{B}_0^f$$

so that $\Gamma^{\text{in}}(0) = 0$ and $\Gamma^{\text{in}}(1) = t^f$. Let $s_1 < s_2 \in (0,1)$ be such that

$$\Gamma^{\text{in}}(s_1) = c^F,\ \Gamma^{\text{in}}(s_2) = p^F.$$

By elementary path-lifting considerations, the curve Γ^{in}_{-1} is a cross-cut in U. Let V be the connected component of $U \setminus \Gamma^{\text{in}}_{-1}$ which does not contain t^F in its boundary. By construction (Proposition 4.10), there exists a univalent branch G of F^{-1} which maps

$$\operatorname{Dom}(F) \setminus \Gamma^{\text{in}}([s_1, 1]) \mapsto V \setminus \Gamma^{\text{in}}([s_1, s_2]).$$

We note:

Proposition 4.13. *The inverse branch G maps the domain $\operatorname{Dom}(F) \setminus \Gamma^{in}$ inside itself.*

Let $l(t)$ be a simple path such that $l(0) = u$, $l(1) = F(u) \in \partial \operatorname{Dom}(F)$, and $\overset{\circ}{l} \cap \overline{U} = \emptyset$. Since $\operatorname{Dom}(F)$ is a Jordan domain, the inverse branch G has a continuous extension to the boundary point u. Set $u_0 \equiv u$, and denote u_{-n} the orbit of u_0 under G. The curve l has a univalent pull-back $l_{-1} \subset \overline{V}$ such that $l_{-1}(0) = u_{-1}$ and $l_{-1}(1) = u$. We let

$$l_{-n} = G(l_{-(n-1)}) \text{ for } n \geq 2.$$

Set

$$\Lambda = \cup_{n\geq 1} l_{-n}.$$

We parametrize this curve by $(0,1]$ so that

$$\Lambda\left(\left[\frac{1}{j},\frac{1}{j-1}\right]\right) = l_{-(j-1)} \text{ for } j \geq 2.$$

Thus, $\Lambda(1/j) = u_{-j}$. By Proposition 4.13 and Denjoy-Wolff Theorem, we have:

Proposition 4.14. *The curve* Λ *is disjoint from* B_0^F*, and lands at* 0*:*

$$\lim_{t\to 0-}\Lambda(t) = 0.$$

Definition 4.4. We call the curve $\hat{\Gamma} \equiv \Gamma \cup \Lambda$ the *secondary* cut.

Let $u^+, u^- \in \partial U$ be the two endpoints of Γ^{in}_{-1}, labeled in such a way that u^+ is encountered first, when going around ∂U in the positive direction from t^f. Denote the connected components of $U \setminus \hat{\Gamma}$ by U^+, U^- so that $U^{\pm} \ni u^{\pm}$.

Consider the inverse branch of f which maps $B_0^F \setminus \Gamma^{\text{in}}([0,s_1])$ into $B_0^F \cap U^+$, and let G_0 be its univalent extension to

$$G_0 : U \setminus (\Lambda \cup \Gamma^{\text{in}}([0,s_1])) \hookrightarrow U^+.$$

Replacing U^+ with U^- we similarly define an analytic branch of F^{-1}

$$G_1 : U \setminus (\Lambda \cup \Gamma^{\text{in}}([0,s_1])) \hookrightarrow U^-.$$

Let $P \subset B_0^F$ be an attracting petal which contains c^F in its interior, and let P_{-1} be a connected component of $F^{-1}(P)$ such that

$$P \subset P_{-1}.$$

We set

$$\mathbf{A} \equiv U \setminus P_{-1}.$$

Let

$$\mathbf{A}_0 \equiv \mathbf{A} \cap U^+ \quad \text{and} \quad \mathbf{A}_1 \equiv \mathbf{A} \cap U^-$$

If $i_0 i_1 \ldots i_n$ is an arbitrary sequence of $n+1$ 0's and 1's, we define

$$\mathbf{A}_{i_0 i_1 \ldots i_n} := \{z \in \mathbf{A}_0 : f^j(z) \in \mathbf{A}_{i_j} \quad \text{for} \quad 1 \leq j \leq n\}$$

The proof of Theorem 4.6 now follows the same lines as in §3.3. It follows from the definitions that

- $\mathbf{A}_{i_0 i_1 \ldots i_n}$ is decreasing in n: $\mathbf{A}_{i_0 \ldots i_n} \subset \mathbf{A}_{i_0 \ldots i_{n-1}}$;
- $F\mathbf{A}_{i_0 i_1 \ldots i_n} = \mathbf{A}_{i_1 \ldots i_n}$;



FIGURE 5. The domain **A**.

- $\mathbf{A}_{i_0 i_1 \ldots i_n} = G_{i_0} \mathbf{A}_{i_1 \ldots i_n}$, and hence
$$\mathbf{A}_{i_0 i_1 \ldots i_n} = G_{i_0} \circ G_{i_1} \circ \cdots \circ G_{i_{n-1}} \mathbf{A}_{i_n},$$
or, more generally,
$$\mathbf{A}_{i_0 i_1 \ldots i_n} = G_{i_0} \circ G_{i_1} \circ \cdots \circ G_{i_{j-1}} \mathbf{A}_{i_j i_{j+1} \ldots i_n} \quad \text{for } 1 \leq j \leq n-1.$$

We show that the diameters of the puzzle pieces $\mathbf{A}_{i_0 i_1 \ldots i_n}$ go to zero as $n \to \infty$, uniformly in $i_0 i_1 \ldots i_n$. More precisely, let us define

$$\rho_n := \sup \left\{ \operatorname{diam}(\mathbf{A}_{i_0 \ldots i_n}) : i_0 \ldots i_n \in \{0,1\}^{n+1} \right\}$$

Since $\mathbf{A}_{i_0 \ldots i_n} \supset \mathbf{A}_{i_0 \ldots i_{n+1}}$, the sequence ρ_n is non-increasing in n, so

$$\rho_* := \lim_{n \to \infty} \rho_n \tag{4.1}$$

exists.

We will prove

Proposition 4.15. *The limit* $\rho_* = 0$.

In the same way as Lemma 3.16, we have

Lemma 4.16. *There is an infinite sequence* $i_0 i_1 \ldots i_n \ldots$ *so that*

$$\operatorname{diam}(A_{i_0 i_1 \ldots i_n}) \geq \rho_* \quad \textit{for all } n.$$

Proof of Proposition 4.15. Let us assume the contrary: $\rho_* > 0$.

We fix a sequence $i_0 i_1 \ldots$ as in the Lemma 3.16, and we split the proof into three cases:

1. i_j is eventually 0;
2. i_j is eventually 1;
3. neither of the above holds.

We start with case (3). There are then infinitely many j's so that

$$i_j = 0 \quad \text{and} \quad i_{j+1} = 1$$

Let j_k be a strictly increasing sequence of such j's. Then, for each k,

$$f_{-j_k} := G_{i_0} \circ G_{i_1} \circ \cdots \circ G_{i_{j_k - 1}}$$

is an analytic branch of the inverse of $f_0^{j_k}$ mapping $\mathbf{A}_{01}$ bijectively to $\mathbf{A}_{i_0 \ldots i_{j_k+1}}$. The closure of $\mathbf{A}_{01}$ does not intersect the postcritical set of f. We now use a version of the argument we gave the proof of Theorem 3.15.

We let Q be an simply connected open neighborhood of $\overline{\mathbf{A}_{01}}$ disjoint from the postcritical set. Then each f_{-j_k} extends to an analytic branch of the inverse of F^{j_k} defined on Q (and we denote this extension also by f_{-j_k}).

By Montel's Theorem, there is a subsequence of (j_k) along which f_{-j_k} converges uniformly on compact subsets of U in the spherical metric on $\hat{\mathbb{C}}$. By adjusting the notation, we can assume that the sequence (f_{-j_k}) itself converges to an analytic function which we denote h. Since

$$\operatorname{diam}(f_{-j_k} \mathbf{A}_{01}) = \operatorname{diam}(\mathbf{A}_{i_0 \ldots i_{j_k+1}})$$

does not go to zero as $k \to \infty$, the function h is non-constant.

Let $z_0 := G_0(u^-) \in \mathbf{A}_{01} \cap \partial B_0^f$. By invarince of the basin boundary, we have $w_0 = h(z_0) \in \partial B_0^f$. Since h is non-constant, $h(\mathbf{A}_{01})$ contains an open neighborhood W of w_0. Let $V \Subset W$ be a smaller open neighborhood of w_0. Then, an arbitrary large iterate F^{j_k} maps V inside Q, which is impossible, since $f(u^-) = t^f \in \partial \operatorname{Dom}(f)$.

We turn next to case 1 above, and deal first with the situation $i_j = 0$ for all j. We write

$$\mathbf{A}^{(n)} := \mathbf{A}_{\underbrace{0 \ldots 0}_{n \text{ terms}}} = (G_0)^{n-2} \mathbf{A}_{00}$$

Now G_0 maps $\mathbf{A}_{00}$ into itself, and $G_0 = G$ on $\mathbf{A}_{00}$, so we can write

$$\mathbf{A}^{(n)} = (G)^{n-2} \mathbf{A}_{00}.$$

By Denjoy-Wolff Theorem applied to g and local dynamics near the parabolic point 0,

$$(G)^{n-2} \to 0 \quad \text{uniformly on } \mathbf{A}_{00},$$

so

$$\operatorname{diam}(\mathbf{A}^{(n)}) \to 0 \quad \text{as } n \to \infty.$$

Next consider sequences of the form $i_0 i_1 \cdots i_k 00 \cdots$, and use the formula

$$\mathbf{A}_{i_0 \ldots i_k \underbrace{0 \cdots 0}_{n \text{ terms}}} = G_{i_0} \circ G_{i_1} \circ \cdots \circ G_{i_k} A^{(n)}.$$

The mapping $G_{i_0} \circ G_{i_1} \circ \cdots \circ G_{i_k}$ extends to be continuous on $\overline{\mathbf{A}_{00}}$, and $\operatorname{diam}(A^{(n)}) \to 0$ by what we just proved, so

$$\operatorname{diam}(\mathbf{A}_{i_0 \ldots i_k \underbrace{0 \cdots 0}_{n \text{ terms}}}) \to 0 \quad \text{as } n \to \infty.$$

A similar argument, using $G_1 = G$ on $\mathbf{A}_{11}$ shows that

$$\operatorname{diam}(\mathbf{A}_{i_0 \ldots i_k \underbrace{1 \cdots 1}_{n \text{ terms}}}) \to 0 \quad \text{as } n \to \infty.$$

□

Let $\mathbf{i} = (i_j)_{j=0}^N$ be a finite or infinite sequence ($N \leq \infty$) of 0's and 1's. We interpret it as a binary representation of a number in $[0, 1]$:

$$\underline{\mathbf{i}}_2 \equiv \sum_{j=0}^{N} i_j 2^j \in [0, 1].$$

For any such dyadic sequence,

$$\overline{\mathbf{A}_{i_0}} \supset \overline{\mathbf{A}_{i_0 i_1}} \supset \cdots \supset \overline{\mathbf{A}_{i_0 \ldots i_n}} \supset \ldots$$

is a nested sequence of compact sets in $\mathbb{C}$ with diameter going to 0, so its intersection contains exactly one point, which we denote by $\hat{z}(\mathbf{i})$. It is immediate from the construction that $\mathbf{i} \mapsto \hat{z}(\mathbf{i})$ is continuous from $\{0, 1\}^{\mathbb{Z}}$ to $\mathbb{C}$. It is not injective, however, this ambiguity is easily tractable:

Lemma 4.17. *Suppose* $\mathbf{i} = i_0 i_1 \ldots i_{n-1}$ *and* $\mathbf{i}' = i'_0 i'_1 \ldots i'_{n-1}$ *are two finite dyadic sequences of an equal length, and*

$$\overline{\mathbf{A}_{\mathbf{i}}} \cap \overline{\mathbf{A}_{\mathbf{i}'}} \neq \emptyset.$$

Then either $\underline{\mathbf{i}}_2 = \underline{\mathbf{i}'}_2$, *or* $\underline{\mathbf{i}}_2 = \underline{\mathbf{i}'}_2 \pm 2^{-n}$.

The proof is a straightforward induction in n, and is left to the reader.

As a corollary, we get:

Corollary 4.18. *Let* $\mathbf{i}$ *and* $\mathbf{i}'$ *be two infinite dyadic sequences. If* $\underline{\mathbf{i}}_2 \neq \underline{\mathbf{i}'}_2$*, then* $\overline{\mathbf{A}_{i_0\cdots i_n}}$ *and* $\overline{\mathbf{A}_{i'_0\cdots i'_n}}$ *are disjoint for large enough* n

Proof. For any $\mathbf{i} = i_0 \cdots i_n \cdots$, and any n

$$|\underline{\mathbf{i}}_2 - \underline{i_0 \cdots i_{n-1}}_2| \leq 2^{-n}$$

so, if n is large enough so that

$$|\underline{\mathbf{i}}_2 - \underline{\mathbf{i}'}_2| > 3 \times 2^{-n},$$

then

$$|\underline{i_0 \cdots i_{n-1}}_2 - \underline{i'_0 \cdots i'_{n-1}}_2| > 2^{-n},$$

which by Lemma 3.17 implies that $\overline{\mathbf{A}_{i_0\cdots i_{n-1}}}$ and $\overline{\mathbf{A}_{i'_0\cdots i'_{n-1}}}$ are disjoint, as asserted. ☐

Putting together what we know about the map $\mathbf{i} \mapsto \hat{z}(\mathbf{i})$, we finally obtain:

Proposition 4.19. *The angles* $\underline{\mathbf{i}}_2 = \underline{\mathbf{i}'}_2$ *if and only if* $\hat{z}(\mathbf{i}) = \hat{z}(\mathbf{i}')$*. Hence there is a function* $\theta \mapsto z(\theta)$ *from the circle* $\mathbb{R}/\mathbb{Z}$ *to* $\mathbb{C}$ *so that*

$$\hat{z}(\mathbf{i}) = \underline{\mathbf{i}}_2.$$

The function $z(\,.\,)$ *is continuous, bijective, and maps the circle onto*

$$J \equiv \overline{\cap_n((F^{-n})|_U \overline{\mathbf{A}})}.$$

By Theorem 3.20, we have:

Corollary 4.20. *The set* J *is a Jordan curve.*

Proposition 4.21. *Let* $\tilde{F} : J \to J$ *be defined as* $\tilde{F}(t^F) = 0$*, and coincide with* F *elsewhere on* J*. The map* $\underline{\mathbf{i}}_2 \mapsto \hat{z}(\mathbf{i})$ *is a conjugacy between* $\theta \mapsto 2\theta$ *and* $\tilde{F}(z)$*:*

$$\tilde{F}(\hat{z}(\mathbf{i})) = \hat{z}(2 \cdot \underline{\mathbf{i}}_2).$$

Let us denote B the connected component of $\hat{\mathbb{C}} \setminus J$ which contains c^F. By construction, $B \cap B_0^F \neq \emptyset$. By the Maximum Principle,

$$F : B \to B.$$

By the classification of dynamics on a hyperbolic domain,

$$B \subset B_0^F.$$

On the other hand, preimages of t^f are dense in $J = \partial B$, and hence

$$B_0^F \cap J = \emptyset.$$

We conclude:

$$B = B_0^F,$$



Figure 6. Several components of the parabolic basin of $f \in \mathbf{H}$

and thus Theorem 4.6 is proven.

We illustrate the topological structure of the parabolic basin of a map $f \in \mathcal{P}(\mathbf{P})$ in Figure 4.3. We have indicated the immediate basin B_0 as well as several components of $f^{-1}(B_0)$. To better understand those, we have drawn the preimage $f^{-1}(C)$ of the circle

$$C = \{|z| = |c|\},$$

which cuts across the only critical value c of f. These critical level curves partition the domain W into univalent preimages of the disks $D^+ = D_{|c|}(0)$ and $D^- = \hat{\mathbb{C}} \setminus \overline{D^+}$.

Two smooth branches in the preimage $f^{-1}(C)$ may intersect at a ramification point of f (all of the ramification points are simple). Each *critical component* of $f^{-1}(B_0)$ has one of these branch points, and touches ∂W in two points ("preimages" of the asymptotic value 0). The *non-critical components* of $f^{-1}(B_0)$ arrange themselves in sequences along the critical level curves, as shown.

4.4. **Convergence of parabolic renormalization.** By definition of the class $\mathbf{P}$, every map $f \in \mathbf{P}$ can be decomposed as

$$f = v \cdot \mathcal{P}(f_0) \circ \varphi_f(e^{2\pi i\theta} z) : \mathrm{Dom}(f) \to \hat{\mathbb{C}},$$

where $v = c^f/c^{\mathcal{P}(f_0)}$. We will denote $\psi_f = \varphi_f^{-1}$. Thus, φ_f conformally maps the unit disk $\mathbb{D}$ onto $W_f = \mathrm{Dom}(f)$ with $\psi_f(0) = 0$ and $\psi'_f(0) > 0$. We will topologize $\mathbf{P}$ by identifying it with the space of thus normalized conformal maps of the unit disk:

$$f \mapsto \psi_f,$$

equipped with the compact-open topology.

Let us state an obvious consequence of the Koebe Distortion Theorem:

Lemma 4.22. *Let $\mathcal{S}$ be a family of univalent maps $h : \mathbb{D} \to \hat{\mathbb{C}}$ with $h(0) = 0$. Assume further that there exist positive constants $0 < a < b$ such that $a \leq |h'(0)| \leq b$ for every $h \in \mathcal{S}$. Then the family $\mathcal{S}$ is equicontinuous.*

H. Inou and M. Shishikura [IS] have recently demonstrated the following:

Theorem 4.23 ([IS])**.** *There exists a class $\mathbf{F}$ of analytic maps with a simple parabolic fixed point at the origin, such that the following properties hold:*

- $\mathcal{P}(\mathbf{F}) \subset (\mathbf{F})$*;*
- *there exists a map $f_* \in \mathbf{F}$ which is a fixed point of the parabolic renormalization:*

$$\mathcal{P}(f_*) = f_*;$$

- *denoting $f_0(z) = z + z^2$, we have $\mathcal{P}(f_0) \in \mathbf{F}$, and*

$$\mathcal{P}^n(f_0) \to f_* \text{ in } \mathbf{F};$$

- *in fact, there is a structure of an infinite-dimensional complex-analytic manifold on $\mathbf{F}$ which is compatible with the local-uniform norm, in which $\mathcal{P}$ is a contraction.*

We remark:

Corollary 4.24. *The fixed point f_* has an analytic extension to a mapping in $\mathbf{P}$ which we will denote in the same way. It is also fixed under $\mathcal{P}$, considered as a transformation $\mathbf{P} \to \mathbf{P}$.*

Proof. By Lemma 4.22, the sequence of parabolic renormalizations $\mathcal{P}^n(f_0)$ is pre-compact in $\mathbf{P}$. By Theorem 4.23, every limit point of this sequence coincides with f_* on a neighborhood of the origin. $\square$

5. NUMERICAL RESULTS

5.1. **Properties of the asymptotic expansion of the Fatou coordinates.** A justification of the use of the asymptotic expansion of the Fatou coordinate (Theorem 2.12) for computation has a theoretical basis in the following. Recall (see e.g. [Ram]) that a formal power series $\sum_{m=1}^{\infty} a_m x^{-m}$ is of *Gevrey order k* if

$$|a_m| < CA^n (n!)^{\frac{1}{k}} \text{ for some choice of positive constants } C, A.$$

As was shown by Écalle [Ec]:

Theorem 5.1. *The asymptotic expansion of Theorem 2.12 is of Gevrey order* 1.

Thus, the asymptotic series of $g(w)$ is rather slowly divergent. A Stirling formula estimate suggest that the first n terms of the series are useful in estimating the value of g for $|w| > \text{const} \cdot n$.

Theorem 5.1 is a part of Écalle's theory of *resurgence* as applied specifically to Fatou coordinates (see [Sau] for an account). Recall, that the *Borel transform* of a formal power series $\sum_{m=1}^{\infty} a_m x^{-m}$ consists in applying the termwise inverse Laplace transform:

$$a_m x^{-m} \mapsto \frac{a_m \zeta^{m-1}}{(m-1)!}.$$

In the case when the formal power series is of Gevrey order 1, this yields a series

$$\sum_{m=1}^{\infty} \frac{a_m \zeta^{m-1}}{(m-1)!},$$

which converges to an analytic function $\hat{h}(\zeta)$ in a neighborhood of the origin. Assume further, that the function $\hat{h}$ extends analytically along the positive real axis, and the extension is of an exponential type. Then, the Laplace transform

$$h(x) = \int_0^{\infty} e^{-x\zeta} \hat{h}(\zeta) d\zeta$$

is defined when $\operatorname{Re} x$ is sufficiently large. The original formal power series $\sum_{m=1}^{\infty} a_m x^{-m}$ is the asymptotic series of $h(x)$, which should thus be considered a summation of the divergent formal series.

Resurgence theory implies much more detailed information about the asymptotic expansion of the Fatou coordinate given in Theorem 2.12, than that required to perform the Borel summation. In particular, the function $\hat{h}$ analytically extends not just along the positive real axis, but along every path in $\mathbb{C} \setminus \{2\pi i n,\ n = \pm 1, \pm 2, \ldots\}$. The attracting Fatou coordinate is obtained using the Laplace transform as described above. The same function $\hat{h}$ also produces the repelling Fatou coordinate: by using the Laplace transform

$$\int_{-\infty}^{0} e^{-x\zeta}\hat{h}(\zeta)d\zeta$$

for x with a large negative real part. The development of the resurgence theory for Fatou coordinates was started by Écalle in [Ec], and was largely completed in the recent work of Dudko and Sauzin [DS].

5.2. **Computational scheme for $\mathcal{P}$.** Having mentioned the resurgent properties of the asymptotic expansion of the Fatou coordinate, we proceed to describe the computational scheme for $\mathcal{P}$. We begin with a germ of an analytic mapping

$$f(z) = z + z^2 + O(z^3)$$

defined in a neighborhood of the origin. Applying the change of coordinates $w = \kappa(z) = -1/z$, we obtain

$$F(w) = w + 1 + \frac{A}{w} + O\left(\frac{1}{w^2}\right)$$

defined in a neighborhood of ∞. We again use the notation $\Phi_A(w)$ for the function which conjugates F with the unit translation

$$\Phi_A(F(w)) = \Phi_A(w) + 1$$

for $\operatorname{Re} w >> 1$. We let $\Phi_R(w)$ be the solution of the same functional equation for $\operatorname{Re} w << -1$. These changes of coordinate are well-defined up to an additive constant, and

$$\phi_A(z) = \kappa^{-1} \circ \Phi_A \circ \kappa(z),\ \phi_R(z) = \kappa^{-1} \circ \Phi_R \circ \kappa(z).$$

As we have seen in Theorem 2.12, the function $\Phi_A(w)$ has an asymptotic development

$$\Phi_A(w) \sim w - A\log w + \mathrm{const}_A + \sum_{k=1}^{\infty} b_k w^{-k}.$$

The coordinate $\Phi_R(w)$ has an *identical* asymptotic development, differing only by the value of const_R. While this may seem surprising at first glance, recall that these functions are Laplace transforms of *different* analytic continuations of the Borel transform of the same divergent series (plus the $w - A\log w + \mathrm{const}$ term).

We select a large integer M (in practice, $M \approx 100$). We will use the asymptotic expansion to estimate $\Phi_A(w)$ for $w \geq M$ and $\Phi_R(w)$ for $w \leq -M$. Consider an iterate $N \approx 2M$ such that

$$\operatorname{Re} F^N(w) \geq M \text{ for } \operatorname{Re} w \in [-M-1, -M].$$

Denote $\nu(z)$ the function

$$\nu(z) = \operatorname{ixp} \circ \Phi_A \circ F^N \circ (\Phi_R)^{-1} \circ \operatorname{ixp}^{-1}(z).$$

It differs from the parabolic renormalization $\mathcal{P}(f)$ only by rescaling the function and its argument:

$$\mathcal{P}(f)(z) = a_1 \nu(a_0 z).$$

Now consider a contour Γ connecting $w = -M - 1 + iH$ with $F(w) \approx -M + iH$ which is mapped onto the circle $S_\rho = \{|z| = \rho\}$ for a small value of ρ by $\operatorname{ixp} \circ \Phi_R$. Select $n \in \mathbb{N}$ and consider the n points in S_ρ given by $x_k = \rho \exp(2\pi k/n),\ k = 0, \ldots, n-1$. We then evaluate the first n coefficients in the Taylor expansion of η at the origin

$$\eta(z) = \sum_{j=0}^{\infty} r_j z^j$$

using a discrete Fourier transform. Specifically, we calculate

$$s_k = \nu(x_k) \approx \sum_{j=0}^{n-1} r_j (x_k)^j = \sum_{j=0}^{n-1} r_j \rho^j \exp(2\pi k j/n),$$

and apply the inverse discrete Fourier transform:

$$r_j \approx \frac{1}{n\rho^j} \sum_{k=0}^{n-1} s_k \exp(-2\pi k j/n).$$

Since

$$\mathcal{P}(f)(z) = \sum_{j=1}^{\infty} s_j a_1 a_0^j z^j,$$

we have

$$a_1 a_0 s_1 = 1, \text{ and further } a_0 = \frac{s_1}{s_2}.$$

This step completes the computation of the Taylor expansion of $\mathcal{P}(f)$.

5.3. **Computing f_*.** In computing the fixed point $f_*(z)$ we find it more convenient to work with the representation of a germ $f(z) = z + z^2 + \cdots$ in the form:

$$f(z) = z \exp(f_{\log}(z)),$$

where $f_{\log}$ is a germ of an analytic function at the origin with $f_{\log}(z) = z + \cdots$. We then rewrite the parabolic renormalization operator in terms of its action on $f_{\log}$:

$$\mathcal{P}_{\log}(f_{\log})(z) = (2\pi i)^{-1}\Phi_A \circ F^N \circ (\Phi_R)^{-1} \circ \mathrm{ixp}^{-1}(z) - \mathrm{ixp}^{-1}(z).$$

This helps to avoid the round-off error which arises from the growth of f_* near the boundary $\partial\,\mathrm{Dom}(f_*)$.

Modifying the scheme described above to the operator $\mathcal{P}_{\log}$, we calculate the fixed point by iterating $\mathcal{P}$ starting at $f_0(z) = z + z^2$:

Empirical Observation 5.2.

$$f_*(z) \approx z + z^2 + 0.(514 - 0.0346i)z^3 + \cdots$$

Our calculations appear reliable up to the size of the round-off error in double-precision arithmetic ($\sim 10^{-14}$) in the disk of radius $r = 5$ around the origin. As we will see below, the true radius of convergence for the series for f_* is approximately 41 (see the Empirical Observation 5.5).

We also estimated the leading eigenvalue of $D\mathcal{P}|_{f_*}$:

Empirical Observation 5.3. *The eigenvalue of $D\mathcal{P}|_{f_*}$ with the largest modulus is*

$$\lambda \approx -0.017 + 0.040i, \ |\lambda| \approx 0.044.$$

The small size of λ explains the rapid convergence of the iterates of $\mathcal{P}$ to the fixed point. To obtain this estimate, we write

$$f(z) = z + z^2 + \sum_{k=3}^{\infty} \mathrm{coeff}_k(f) z^k,$$

and consider the spectrum of the $N \times N$ matrix $A = (a_{ij})_{i,j=3..N+3}$, with

$$a_{ij} = \frac{\mathrm{coeff}_j(\mathcal{P}(f_* + \epsilon z^i)) - \mathrm{coeff}_j(f_*)}{\epsilon},$$

which serves as a finite-dimensional approximation to $D\mathcal{P}|_{f_*}$.



FIGURE 7. The domain of analyticity of $\mathcal{P}f_0(z)$ for $f_0(z) = z + z^2$, with the immediate parabolic basin indicated.

5.4. **Computing a tail of the domain Dom(f_*).**

Computing the tail using an approximate self-similarity near the tip. Let us denote

$$t_* \equiv t^{f_*} = \partial \mathrm{Dom}(f_*) \cap \overline{B_0^{f_*}}$$

the endpoint of the tail of the immediate basin of f_*. Let C_R be a repelling fundamental crescent of f_*, and let $w \in C_R$ have the property

$$t_* = \mathrm{ixp} \circ \phi_R(w).$$

Let $k \geq 2$ be such that

$$f_*^k(w) = 0, \text{ so that } f_*^{k-1}(w) = t_*.$$

Denote χ the local branch of $f_*^{-(k-1)}$ which sends t_* to w. Then the composition

$$\nu \equiv \mathrm{ixp} \circ \phi_R \circ \chi$$

is an analytic map defined in a neighborhood of the endpoint t_*, which fixes it:

$$\nu(t_*) = t_*.$$

This point can be found numerically:

Empirical Observation 5.4.

$$t_* \approx -779.306 - 643.282i, \text{ and } \nu'(t_*) \approx 0.232 + 0.264i.$$

Thus, we have identified the endpoint of the largest tail of Dom(f_*). This construction also gives us the means to compute the tail itself. This can be done by successively applying ν to the immediate basin $B_0^{f_*}$ thus pulling it in towards t_*.

Now let $q \in C_R$ be any other preimage of 0:

$$f_*^l(q) = t_* \text{ for some } q \in \mathbb{N}.$$

Then $v = \mathrm{ixp} \circ \phi_R(q)$ is the endpoint of a different tail in $\partial\mathrm{Dom}(f_*)$. It can be computed by first pulling back the tail of $B_0^{f_*}$ using the inverse branch

$$f_*^{-l} : t_* \mapsto q,$$

and then applying $\mathrm{ixp} \circ \phi_R$.

Computing the tail using the functional equation for an inverse branch. A more careful analysis of the tail can be done as follows. Denote ξ the local branch of f_*^{-1} defined in a slit neighborhood $D_r(0) \setminus [0, r)$ for some small value of r, which sends $0 \mapsto t_*$. We


χ
B_0
u
w
C_r
ξ
$\text{ixp} \circ \tilde{\phi}_R$
Dom(f_*)
w
$\text{ixp} \circ \tilde{\phi}_A$

Figure 8. The inverse branches used in computing the tail of Dom(f_*).

can write the renormalization fixed point equation for this particular branch:

$$\xi = \psi_R \circ \xi \circ \psi_A^{-1}, \tag{5.1}$$

where $\psi_R = \chi \circ \text{ixp} \circ \phi_R$, and ψ_A^{-1} is the appropriately chosen branch of $(\text{ixp} \circ \phi_A)^{-1}$ (thus the "self-similarity" of the tail is exponential, rather than linear). We are going to use the renormalization equation (5.1) *inductively* to compute $\xi(z)$ for sufficiently small values of z, and thus plot the tail.

Representing the numbers in the image of the tail. Numerical computations indicate that the value of $r = 0.0002$ is sufficiently small for our needs, and for $|z| < r$ the difference between the left and the right sides of (5.1) is of the order of 10^{-11}. The values of z for which we would like to evaluate $\xi(z)$ become too small to be represented by the standard double precision numbers (and even too small for their logarithms to be so represented). We write

$$s(t) = \exp(2\pi t),$$

and choose $\hat{t}$ so that

$$\exp(-2\pi\hat{t}) = 0.0002, \text{ that is } \hat{t} = 1.3555...$$

We then represent a small positive number x as

$$x = \frac{1}{s^k(t)},$$

for the unique choices of $t \in [\hat{t}, s(\hat{t}))$, and an iterate $k \in \mathbb{N}$.

We can write any complex number z with $|z| < r$ uniquely as

$$z = (k, t, \theta) \equiv \frac{\exp(2\pi i\theta)}{s^k(t)}, \ 0 \leq \theta < 1.$$

Note that this representation of small numbers makes it very easy to compute logarithms. In particular,

$$\text{ixp}^{-1}((k, t, \theta)) = \theta + is^{k-1}(t).$$

The next step in applying (5.1) is to apply ϕ_A^{-1} to the right-hand side of the equation. From the first two terms in the asymptotics of

$$\phi_A(z) = -\frac{1}{z} + O(\log|z|) \text{ for small } z,$$

it follows that

$$\phi_A^{-1}(y) = -\frac{1}{y + O(\log|y|)} \text{ for large } |y|.$$

A numerical estimate shows that for $|y| \geq 10^{18}$ the $O(\log|y|)$ term dissapears into the round-off error, when added to y. Thus

$$\psi_A^{-1}((k, t, \theta)) \approx -\frac{1}{\theta + is^{k-1}(t)} \approx is^{k-1}(t) = (k-1, t, 1/4)$$

provided $s^{k-1}(t) \geq 10^{18}$. A direct estimate shows that for either $k \geq 3$, or $k = 2$ and $t > 18\log 10/2\pi \approx 6.596$ the last inequality will hold.

The size of the domain of analyticity. To draw the pictures of the domain of analyticity of the fixed point of f_* (Figures 5.4 and 5.4) we employed the following strategy. First, a periodic orbit of period 2 in ∂B_0 was identified. Its preimages give a rough outline of ∂B_0, but become sparse near the “tails”, which are not visible in this initial outline. At the next step, the large “tail” of B_0 is computed as described above. Finally, its preimages are used to fill in the remaining gaps in ∂B_0.



Figure 9. The domain of analyticity of f_* and the boundary of the immediate parabolic basin $B_0^{f_*}$. In the second figure, a part of the critical level curve of f_* is also indicated.



FIGURE 10. A blow-up of the boundary of the immediate basin of f_* in the vicinity of the parabolic point.

As the final step, we calculate the boundary of $\mathrm{Dom}(f_*)$ as

$$\partial\,\mathrm{Dom}(f_*) = \mathrm{ixp} \circ \phi_R(\partial B_0 \cap P_R).$$

An empirical estimate of the inner radius of $\mathrm{Dom}(f_*)$ around the origin allows us to formulate the following observation (see Figure 5.4):

Empirical Observation 5.5. *The radius of convergence of the Taylor's expansion of* f_* *at the origin is* $R \approx 41$.

References

[BE] X. Buff, A. Epstein. A parabolic Pommerenke-Levin-Yoccoz inequality. *Fund. Math.* 172, no 3.(2002), 249-289

[Do1] A. Douady, *Systéms dynamiques holomorphes*, Seminar Bourbaki, Astérisque, **105-106** (1983) 39-64.

[Do2] A. Douady. Does a Julia set depend continuously on the polynomial? In *Complex dynamical systems: The mathematics behind the Mandelbrot set and Julia sets.* ed. R.L. Devaney, Proc. of Symposia in Applied Math., Vol 49, Amer. Math. Soc., 1994, pp. 91-138.

[DH] A. Douady, J.H. Hubbard. Etude dynamique des polynômes complexes, I-II. Pub. Math. d'Orsay, 1984.

[DS] A. Dudko, D. Sauzin. Manuscript in preparation, 2011.

[Ec] J. Écalle. Les fonctions résurgentes, Publ. Math. d'Orsay [Vol. 1: 81-05, Vol. 2: 81-06, Vol. 3: 85-05] 1981, 1985

[Ep] A. Epstein, Towers of finite type complex analytic maps. PhD Thesis, CUNY, 1993.

[EKT] A. Epstein, L. Keen, C. Tresser. The set of maps $F_{a,b} : x \mapsto x + a + \frac{b}{2\pi}\sin(2\pi x)$ with any given rotation interval is contractible. *Commun. Math. Phys.* **173**, 313-333, 1995.

[EY] A. Epstein, M. Yampolsky. The universal parabolic map. Accepted to *Ergodic Th. & Dynam. Sys.*, available from *http://www.math.toronto.edu/yampol*

[IS] H. Inou and M. Shishikura. The renormalization for parabolic fixed points and their perturbation. Preprint, 2008.

[Lan] O.E. Lanford. The zero rotation number fixed point. Manuscript.

[Lav] P. Lavaurs. Systémes dynamiques holomorphes: Explosion de points périodiques, Thèse, Université de Paris-Sud, 1989.

[Lyu] M. Lyubich. The dynamics of rational transforms: the topological picture. Russian Math. Surveys 41:4(1986), 43-117.

[Mil1] J. Milnor. Dynamics in One Complex Variable. 3rd ed. Annals of Math. Studies 160. Princeton University Press, 2006

[Mil2] J. Milnor, Pasting together Julia sets - a worked out example of mating, Experiment. Math. **13**(2004), no. 1, 55–92.

[Ram] J.-P. Ramis. Séries divergentes et théories asymptotiques, Panoramas et Synthèses (1994)

[Sau] D. Sauzin. Resurgent functions and splitting problems. RIMS Kokyuroku 1493, p. 48-117, 2005 available from *http://www.imcce.fr/Equipes/ASD/person/Sauzin/sauz_preprint.php*

[Sh] M. Shishikura. The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets. *Ann. of Math.* (2) **147**(1998), no. 2, 225-267.

[Vor] S.M. Voronin. Analytic classification of germs of conformal maps $(\mathbb{C}, 0) \to (\mathbb{C}, 0)$ with identity linear part. *Funct. Anal. Appl.*, **15**(1981), 1-17.

[Ya] M. Yampolsky. The attractor of renormalization and rigidity of towers of critical circle maps, *Commun. Math. Phys.*, **218**(2001), no. 3, 537-568.

[YZ] M. Yampolsky, S. Zakeri. Mating Siegel quadratic polynomials, *Journ. Amer. Math. Soc.*, **14**(2000), 25-78